PROFICIENCY TESTING IN BONE MARROW MORPHOLOGY

John Sioufi, Katherine Marsden
RCPAQAP Haematology, St Leonards, NSW, Australia

Aim: The RCPAQAP blood film morphology module originated in 1963 unable to offer a bone marrow (BM) morphology module as it was not possible to obtain the number of slides required for a survey. Virtual microscopy overcomes this problem and the aim was to determine the feasibility of a BM morphology survey using online virtual microscope scans.

Method: A pilot survey was sent to all laboratories enrolled in the RCPAQAP Morphology module. Virtual microscopy scans of the blood film, bone marrow aspirate and trephine from an essential thrombocythaemia (ET) patient were used. Participation involved providing a bone marrow differential count, a diagnosis and recommended additional investigations.

Results: The uptake was low despite reminders and extensions of the closing date. Forty participants completed the differential count; 30–39 answered survey questions and/or provided other feedback. The diagnoses made were ET 31/39 participants (80%), other myeloproliferative neoplasms 5/39 (13%), other diagnoses 3 (8%).

Conclusion: The reasons for the low response rate are not known but we conclude that many laboratories could now manage an online virtual bone marrow morphology survey. As a result the RCPAQAP will introduce a bone marrow morphology module in 2016 but ensure that DVDs are available if preferred.

THROMBOPHILIA INVESTIGATION – NOT FOR EVERYONE

Sam Schulman
Department of Medicine, McMaster University, Hamilton, ON, Canada

‘All truths are easy to understand once they are discovered; the point is to discover them’ (Galileo Galilei). Many physicians believe that a thrombophilia work-up will provide the explanation for a thrombotic event and thousands of dollars have been spent on these investigations. For arterial thromboembolic events a thrombophilia investigation is rarely positive and it will never lead to any beneficial change in management of the patients. In venous thromboembolism it can demonstrate abnormalities in up to 50% of the cases, depending on how they are selected. False positive results are common if the blood sampling is performed when an anticoagulant drug is present. The most affected tests are lupus anticoagulant, protein C, protein S and antithrombin. The acute phase reaction as well as liver dysfunction secondary to large pulmonary embolism with right heart strain can also distort the results. A negative investigation does not mean that the patient has no thrombophilic condition, since some abnormalities surely remain to be discovered. Before embarking on a thrombophilia work-up some questions need to be answered: (1) will the result change the management—probably not if the condition requires indefinite anticoagulation—and (2) if not, is it for information and preventive measures for family members, and in that case, are they interested? The evidence that such information will change prognosis is very weak.

NEW ANTICOAGULANTS, WHY ARE THEY DIFFERENT

Sam Schulman
Department of Medicine, McMaster University, Hamilton, ON, Canada

The non-vitamin K antagonist oral anticoagulants (NOACs) have several characteristics that differ from vitamin K antagonists (VKAs). NOACs have quick onset and shorter half-life, which are mostly favourable features. However, for patients with poor adherence the short half-life poses a greater risk of lost efficacy, together with the absence of supervision required for the VKAs. The risk of bleeding is similar or lower with NOACs; most importantly, there is a reduction of the risk of intracranial haemorrhages, explained by the severe impairment of tissue factor pathway initiation of coagulation with VKAs but not with NOACs. On the other hand, NOACs are associated with more bleeding from the distal gastrointestinal canal due to the high concentration of active drug in the bowel. This is a result of the efflux transporter, P-glycoprotein. A similar mechanism may be responsible for the increased incidence of menorrhagia with some NOACs. The management of anticoagulants for surgery must take into account the different pharmacokinetic characteristics of the anticoagulants. Local protocols should be established to standardise peri-operative management in order to minimise the risk for thromboembolic and haemorrhagic complications. The NOACs have more or less dependence on renal function for elimination, which limits their use. Drug-drug interactions are substantially fewer for the NOACs. Rivaroxaban at treatment doses is dependent on concomitant food intake for optimal absorption.

B-CELLS BEHAVING BADLY I: A BETTER BASIS TO BEHOLD BELLIGERENCE IN SMALL B-CELL LYMPHOMAS

Adam Bagg
Department of Pathology and Laboratory Medicine, University of Pennsylvania, Philadelphia, PA, USA

Small B-cell lymphomas and leukaemias (SBCLs) are a clinically, morphologically, immunophenotypically and genetically heterogeneous group of neoplasms that include chronic lymphocytic leukaemia/small lymphocytic lymphoma (CLL/SLL), mantle cell lymphoma (MCL), follicular lymphoma (FL), lymphoplasmacytic lymphoma (LPL), marginal zone lymphoma (MZL) and hairy cell leukaemia (HCL). The pathogenesis of some of these lymphoid malignancies is characterised by distinct translocations, for example t(11;14) and t(14;18), respectively, in most cases of MCL and FL. By contrast, other entities are associated with a variety of non-specific and non-diagnostic but nevertheless recurrent and prognostically pertinent numeric abnormalities, such as del(17p), del(13q14), del(11q22) and +12 in CLL. Yet others like LPL and HCL lack specific cytogenetic aberrations. The recent surge in next generation sequencing technology has shed abundant light on the genetic landscape of SBCLs through characterisation of numerous driver mutations including *BRAF* (V600E) in HCL and *MYD88* (L265P) in LPL, amongst others. These significant advances and panoply of available tests notwithstanding, it is important to appreciate that not all SBCLs need genetic testing for diagnosis, not all harbour their characteristic genetic features