showed a polyp composed of numerous ductal structures lined by a double layer of cells comprising an inner layer of bland ductal epithelium and an outer layer of myoepithelial cells.

To the best of our knowledge twelve cases of epithelial myoepithelial carcinoma of the sinonasal region have been reported in the literature, including the present case. In review of the literature, most cases, including the case presented, were disease free at the 10–40 month post-operative follow up. This neoplasm is a typically a low grade malignant neoplasm, but in rare cases the tumour can metastasise to distant sites.

In conclusion, we illustrate a rare case of epithelial-myoepithelial carcinoma of the sinonasal cavity, a morphologically characteristic tumour in an unusual site.

References

1. Barnes L, Eveson JW, Reichart P, et al. World Health Organization Classification of Tumours: Pathology and Genetics of Head and Neck Tumours. Lyon: IARC Press, 2005; 225–6.
2. Thompson L, Wenig B. Diagnostic Pathology: Head and Neck. Salt Lake City: Amirsys, 2011; 5-112–7.
3. Seethala RR, Barnes EL, Hunt JL. Epithelial myoepithelial carcinoma: A review of the clinicopathologic spectrum and immunophenotypic characteristics in 61 tumours of the salivary gland. *Am J Surg Pathol* 2007; 31: 44–57.

A RARE VASCULAR TUMOUR OF THE OVARY: A CASE REPORT

Yeu Sheng Ang, Gregory Imhagwe, Denys Fortune
Dorevitch Pathology, Traralgon, Vic, Australia

Case history: A 72-year-old female presented with a 2 year history of a right adnexal mass. Right salpingo-oophorectomy revealed a 40 mm diameter, circumscribed mass with a spongy, haemorrhagic cut surface. Microscopically, the mass was composed of small to medium sized blood vessels within an oedematous and fibrotic stroma, with islands of anastomosing vascular channels lined by atypical endothelial cells showing moderate nuclear pleomorphism and occasional mitotic figures. Focally within some of the anastomosing vascular channels a few haematopoietic cells were seen. The endothelial cells were positive for vascular markers CD31 and CD34. Small nests of luteinised stromal cells, showing inhibin positivity were noted at the periphery of the tumour, consistent with residual ovarian parenchyma.

Discussion: We present this as a case of primary ovarian angiosarcoma. Primary ovarian angiosarcoma is a rare entity. Angiosarcomas may display a wide range of morphological features ranging from well (low grade) to poorly (high grade) differentiated. Poorly differentiated lesions show areas of sarcomatous differentiation with significant nuclear atypia, brisk mitotic activity, necrosis and destructive growth. Differentiating between benign vascular tumours and well-differentiated angiosarcomas may be challenging.

GENOMICS VS PLASIAS: GENETIC EXPLANATIONS FOR THE HISTOPATHOLOGIC FEATURES OF TUMOURS

Leon P. Bignold

Division of Surgical Pathology, SA Pathology, Adelaide, SA, Australia

Tumours result from genomic disturbances in somatic cells. In genetics, genotypic changes are called ‘mutations’ and phenotypic changes are called abnormal ‘traits’. However, genetic terms are not used for the histopathology of tumours. Instead, 19th century words such as ‘(de)differentiation’, ‘neoplasia’, ‘dysplasia’ and ‘anaplasia’¹ are employed.

Nevertheless, histopathologic phenomena of tumours can be analysed in genetic terms.² First, tumours generally exhibit excessive growth together with other traits, particularly altered specialisation, abnormal cellular spatial arrangements, and invasion and metastasis. The phenomenon of association of traits is known as ‘linkage’. Second, linkage of traits is usually a manifestation of proximity of relevant genes in complex loci. Hence third, if in a somatic cell mutation causes linked traits, the relevant genomic lesion probably involves a complex locus. That is: a larger lesion than a single gene, or in a particular nucleotide in an exon of a particular gene. Several mechanisms for mutations of entire complex loci are known.

Lastly, additional phenomena of tumour types, for example their specificity to kinds of parent cell, and their variabilities, can be explained by genomic mechanisms.

References

1. Bignold LP, Coghlan BLD, Jersmann HPA. *David Paul Hanseman, Contributions to Oncology – Context, Comments and Translations*. Basel: Birkhäuser, 2007.
2. Bignold LP. *Principles of Tumors*. Waltham: Academic Press/Elsevier, 2015; 222–32.

METASTATIC THYMOMA – A RARE PRESENTATION AS A DURAL-BASED SPINDLE CELL LESION IN THE CRANIAL CAVITY

Manish Mahadeorao Bundelee, Lee Hwei Yee

Department of Pathology, Tan Tock Seng Hospital, Singapore

Intracranial dural-based spindle cell lesions are most commonly meningiomas, solitary fibrous tumours and haemangiopericytomas. Other primary and metastatic spindle lesions are rare. Metastatic thymoma presenting as an intracranial lesion is exceptional and poses a diagnostic challenge.

We report the case of a 75-year-old Indonesian female who presented with a solitary 2.8 cm dural-based mass in the left parieto-occipital region. Excision showed a cellular spindle cell tumour with a prominent haemangiopericytoma-like vascular pattern. Frequent rosettes were seen. The tumour cells showed bland bipolar nuclei, fine chromatin, inconspicuous nucleoli and rare mitoses. Dispersed small lymphocytes were present. The spindled tumour cells were positive for cytokeratins AE1/3 and Cam5.2, PAX8 and CD20. Terminal deoxynucleotidyl transferase (TdT) and CD5 showed immature T-lymphocytes within the tumour. The spindle cells were negative for synaptophysin, chromogranin, epithelial membrane antigen, CD34, and TTF1. The features were highly suggestive of metastatic thymoma with Type A or Type AB features. Computed tomography showed a large anterior mediastinal mass, consistent with a primary mediastinal thymoma.

Thymic epithelial tumours with intracranial metastases are unusual and most are thymic carcinomas. Thymomas with intracranial metastases are rare. The characteristic morphology and immunoperoxidase profile of thymomas are of value in the differential diagnostic work-up.