¹General Medicine, ²Endocrine and Metabolic Units, Royal Adelaide Hospital, ³Discipline of Medicine, University of Adelaide, and ⁴Chemical Pathology Directorate, SA Pathology, Adelaide, SA, Australia

Non-islet cell tumour-induced hypoglycaemia (NICTH) is a rare but important cause of hypoglycaemia. Here we report a case of NICTH secondary to progressive metastatic haemangiopericytoma.

A 39-year-old man was found unresponsive, with low blood glucose recorded: he responded well to intramuscular glucagon and oral glucose. His history was significant for past resection of a primary meningeal haemangiopericytoma, and known hepatic metastases. He had no history of diabetes or access to hypoglycaemic agents. There was no evidence of infection. Thyroid function and serum cortisol were within normal limits.

Subsequent observation showed recurrent episodes of hypoglycaemia. Samples taken during an episode showed a blood glucose concentration of 3.1 mmol/L [reference interval (RI): 3.2–5.5], and suppressed serum insulin [0.5 mU/L (RI: 0–12)], C-peptide [<100 pmol/L (RI: 300–1600)] and insulin-like growth factor (IGF)-1 [4 nmol/L (RI: 10–28)] levels. IGF-2 was not suppressed [72.1 nmol/L (RI: 47–94)]. Imaging revealed progression of metastatic disease.

The combination of low insulin, C-peptide and IGF-I and inappropriately elevated IGF-2 in the presence of low plasma glucose is consistent with tumoural IGF-2 secretion and subsequent NICTH, probably related to increased tumour burden.

Clinicians should be aware of this potentially life-threatening paraneoplastic syndrome.

NON-KETOTIC HYPERGLYCINAEMIA IN A NEONATE DIAGNOSED BY MULTIDISCIPLINARY APPROACH

Geetha Rathnayake, Kay Weng Choy, James C.G. Doery
Monash Pathology, Monash Medical Centre and Department of Medicine, Monash University, Clayton, Vic, Australia

Case presentation: A 5-day old term baby boy, after elective caesarean delivery and APGARs (9/9), presented with lethargy, poor feeding, twitching and seizure activity in all limbs. He was afebrile and never cried.

Results: Electrolytes, CRP, lactate, ammonia, pyruvate and capillary blood gases were normal. CSF total protein was 1.7 g/L (RR 0.1–0.3) with normal glucose, lactate, and microscopy with no growth on culture. CSF viral studies were normal. FBE showed no evidence of infection. EEG indicated encephalopathy. MRI diffusely abnormal and MRI spectroscopy showed possible glycine peak. Subsequent CSF analysis showed elevated glycine of 105 μ mol/L (RR 5–8). Plasma amino acids also showed very high glycine of 862 μ mol/L (RR 106–254). Glycine was also elevated in the urine metabolic screening.

Discussion: The more common causes of encephalopathy were excluded (neonatal hypoxia, infection, infarction, haemorrhage) while MRI spectroscopy suggested possible metabolic disorder. Elevated glycine in blood, urine and especially CSF confirmed a diagnosis of autosomal recessive non-ketotic hyperglycinaemia due to a metabolic block in the mitochondrial glycine cleavage system which converts glycine to CO₂, NH₃ and hydroxymethyltetrahydrofolic acid.

Anticonvulsants and sodium benzoate reduced seizures, improved feeding and level of consciousness. Prognosis however remains poor.

Conclusion: A multidisciplinary approach optimises diagnosis of metabolic disorders.

DIAGNOSIS OF WILSON'S DISEASE – A 20 YEAR AUDIT

Geetha Rathnayake¹, Mirette Saad², Kay Weng Choy¹, James C.G. Doery^{1,3}

¹Monash Pathology, Monash Medical Centre, ²Healthscope Pathology, and ³Department of Medicine, Monash University, Clayton, Vic, Australia

Aims: To review the clinical presentation and relative value of pathology tests in the diagnosis of Wilson's disease.

Methods: Records of all patients diagnosed with Wilson's disease in the hospital medical record or pathology information system were reviewed.

Results: A total of 18 patients were diagnosed over a 20-year period and were aged 10–54 years; 78% were male. Based on surname 55% appeared to be of Asian origin. The majority (67%) were 10–20 years, of whom 90% presented with liver disease. Of those over 20 years 83% had neuro-psychiatric symptoms.

No single biochemistry test had 100% sensitivity. Thus serum copper was low in 86%, caeruloplasmin low in 86% while urine copper was high in 85%. While no patients had a normal serum copper, surprisingly two had elevated levels in the setting of acute/sub-acute hepatitis. Liver copper was high in all patients tested ($n=6$).

Conclusion: While no blood or urine test demonstrated 100% sensitivity, all patients had at least one of the key tests outside the reference range. This audit confirms the general literature on Wilson's disease and underscores the need for Chemical Pathologists and other clinicians to appreciate the limitation of the individual tests and to communicate effectively to ensure accurate diagnosis.

IMPROVING INTERPRETATION OF FLAT ORAL GLUCOSE TOLERANCE TEST IN PREGNANCY

M.M. Salib¹, A.J. Simpson¹, P.E. Hickman^{1,2}, S. Apostoloska¹, C. Yu¹, J.M. Potter^{1,2}

¹ACT Pathology, The Canberra Hospital, Canberra, and ²ANU Medical School, Australian National University, Acton, ACT, Australia

Aim: Gestational diabetes mellitus has a strong association with adverse pregnancy outcomes. The oral glucose tolerance test (OGTT) is the 'gold standard' for its diagnosis. A flat OGTT is a significant problem encountered with interpreting results. To interpret the flat curve, we implemented testing plasma insulin.

Methods: Over 8 months, 330 flat OGTT have been identified (11% of all pregnant women using the 75 g 2 h OGTT). Data were analysed according to the insulin response and the difference in glucose concentration between 1 and 2 h.