11. Uhm J, Hamad N, Michelis FV, *et al.* The risk of polyomavirus BK-associated hemorrhagic cystitis after allogeneic hematopoietic SCT is associated with myeloablative conditioning, CMV viremia and severe acute GVHD. *Bone Marrow Transplant* 2014; 49: 1528–34.
12. Jacobi J, Prignitz A, Büttner M, *et al.* BK viremia and polyomavirus nephropathy in 352 kidney transplants; risk factors and potential role of mTOR inhibition. *BMC Nephrol* 2013; 14: 207.

DOI: <https://doi.org/10.1016/j.pathol.2019.05.007>

Development of monomorphic B-cell plasmacytoma post-transplant lymphoproliferative disorder in a patient with previous autologous stem cell transplant

Sir,

Post-transplant lymphoproliferative disorders (PTLDs) are uncommon lymphoid and/or plasmacytic proliferations that can occur in patients after undergoing solid organ or haematopoietic stem cell transplantation, secondary to extrinsic immunosuppression inherent in these procedures. In the context of haematopoietic stem cell transplantation, the large majority of reported PTLDs occur in patients who received allogeneic haematopoietic stem cell transplantation (AlloHSCT). Only a few cases of PTLD following autologous haematopoietic stem cell transplantation (AHSCT) have been reported.^{1–6} We herein report the rare interesting case of an Epstein–Barr virus (EBV) negative monomorphic B-cell plasmacytoma post-transplant lymphoproliferative disorder (PTLD), developing in a patient who received AHSCT for treatment of advanced stage classic Hodgkin lymphoma (CHL).

A 23-year-old male presented to our institution 4 years previously with fever, weight loss and shortness of breath of 2–4 months duration. Positron emission tomography (PET) scan performed showed a fluorodeoxyglucose (FDG) avid, large, bulky, 10.6 cm anterior mediastinal mass, multiple FDG avid lymph nodes above and below the diaphragm, as well as multiple FDG lesions in the axial skeleton. A biopsy of the anterior mediastinal mass was performed, and the patient was diagnosed with nodular sclerosis CHL (Fig. 1A). The patient was treated with six cycles of adriamycin, bleomycin, vinblastine and dacarbazine (ABVD) and radiotherapy to the mediastinum. The patient achieved complete remission based on positron emission tomography (PET) scan findings after the treatment. However, 2 years later he developed new FDG avid lymphadenopathy above and below the diaphragm, as well as new FDG avid iliac bone lesions. Biopsy of the iliac bone showed disease relapse of CHL. The patient received platinum-based salvage chemotherapy for the disease relapse. After harvesting of peripheral blood stem cells, he underwent AHSCT. The conditioning regimen consisted of carmustine, etoposide, cytarabine and melphalan. Unmanipulated haematopoietic stem cells were administered at a cell dose of 2.78×10^6 CD34⁺ cells/kg. The patient had an episode of culture negative neutropenic fever post-AHSCT, from which he subsequently recovered. He was not put on any immunosuppressive drug regimen post-AHSCT. Three and a half months following AHSCT, the follow-up PET scan showed new FDG avid left submental and level II cervical lymph nodes. An excision biopsy of a left level II cervical lymph node was performed. On histology, the cervical lymph node architecture was preserved with presence of reactive lymphoid follicles. There were increased numbers of small mature plasma cells noted in the paracortical areas and sinuses (Fig. 1B). No neoplastic Reed–Sternberg cells were seen. The histological diagnosis

Fig. 1 (A) First mediastinal mass biopsy showing neoplastic classical Reed–Sternberg cells surrounded by reactive inflammatory cells (H&E). (B) Subsequent left level II cervical lymph node biopsy showing preserved nodal architecture, but with increased numbers of plasma cells in the paracortical areas (H&E). (C,D) Final left submental lymph node biopsy showing effacement of nodal architecture by sheets of mature plasma cells and plasmablasts (H&E).

at that time was a lymph node with reactive changes. However, the patient had a persistently enlarged submental lymph node on follow-up, which was excised 2 months later. On histology, the submental lymph node showed notable architectural effacement by confluent sheets of small mature plasma cells and medium to large lymphoid cells with eccentric nuclei and conspicuous nucleoli, consistent with plasmablasts (Fig. 1C,D). Immunohistochemical (IHC) studies were performed (Fig. 2A–G). The plasma cells and plasmablasts were positive for CD79A and CD138. Only a small proportion of plasmablasts expressed CD20 and CD30. The plasma cells and plasmablasts were negative for CD3, PAX5 and CD56. Kappa and lambda light chains encoded RNA *in situ* hybridisation was performed. The plasma cells and plasmablasts showed evidence of lambda light chain restriction, indicating a monoclonal mature B-cell population. EBV encoded RNA (EBER) *in situ* hybridisation was also performed. The plasma cells and plasmablasts showed no evidence of EBV infection. The Ki-67 proliferative index of the plasmacytic/plasmablastic infiltrate was up to ~50%. Immunoglobulin heavy chain (IgH) gene rearrangement study performed using BIOMED2 primers showed a

monoclonal gene rearrangement (Fig. 2H). A final diagnosis of an EBV-negative monomorphic B-cell plasmacytoma PTL was made. Bone marrow studies, bone skeletal survey and other blood investigations performed excluded the diagnosis of myeloma. A subsequent PET scan showed localised disease, comprising mild uptake FDG-avid small volume lymph nodes, confined to the cervical region. The patient was otherwise clinically well with no B symptoms of fever, night sweats and weight loss. After a multi-disciplinary tumour board discussion, the decision was made for watchful waiting with no chemotherapy given for his PTL. Three months later, follow-up PET scan surprisingly showed resolution of FDG-avid lymph nodes with minimal uptake seen. The resolution was attributed to the patient's immune reconstitution.

PTLDs are lymphoid or plasmacytic proliferations that can occur as a result of immunosuppression after solid organ and haematopoietic stem cell transplant (HSCT). The four main categories of PTLDs are non-destructive (early), polymorphic, monomorphic and classic Hodgkin lymphoma.⁷ PTLDs are more often EBV positive, but can also be EBV negative. In contrast to EBV positive PTLDs,

Fig. 2 (A) The plasma cells and plasmablasts are positive for CD79A and (B) CD138. (C) The Ki-67 proliferative index is up to ~50%. (D) Kappa-ISH and (E) lambda-ISH: the plasma cells and plasmablasts show evidence of lambda light chain restriction. (F) The plasma cells and plasmablasts are negative for EBER-ISH and (G) CD56. (H) A monoclonal gene rearrangement is detected on immunoglobulin heavy chain (IgH) gene clonality assay by polymerase chain reaction.

EBV negative cases are more common in adults, of later onset after transplant, and are more likely to be of monomorphic subtype.⁸ Most cases of PTLTD developing after HSCT occurred in the first 6 months post-transplant.⁹ There have only been a few reported cases, mostly in the form of case reports, of PTLTDs developing after AHSCT as compared to AlloHSCT. This is unsurprising as AHSCT involves a lesser degree of immune suppression than AlloHSCT. The large majority of reported cases of PTLTDs developing after AHSCT are EBV positive. Only one case of an EBV negative PTLTD with features of Hodgkin lymphoma has been reported.³ To the best of our knowledge, ours is the first reported case of an EBV negative monomorphic B-cell plasmacytoma PTLTD following AHSCT. The main differential diagnosis considered for our case was a *de novo* primary nodal plasma cell neoplasm/plasmacytoma. The diagnosis of PTLTD, rather than a *de novo* plasmacytoma, was supported by disease presentation within 6 months following AHSCT and the apparent improvement of disease without chemotherapy. The disease improvement was attributed to the gradual full restoration of the patient's immune system after the immunosuppressive effects of the AHSCT had dissipated. The histological findings of plasmacytic hyperplasia with preserved nodal architecture in the initial cervical lymph node biopsy specimen were most likely on hindsight a form of non-destructive PTLTD. This was followed by progression into a monomorphic PTLTD with architectural effacement, as evident in the subsequent submental lymph node biopsy specimen. Non-destructive PTLTDs, in contrast with the other subtypes, are defined as lymphoid or plasmacytic proliferations that do not show architectural effacement and do not meet any diagnostic criteria of any known lymphoid neoplasms. The old term of 'early PTLTD' was dropped in the revised 4th Edition World Health Organization (WHO) classification to avoid confusion with various PTLTDs that occur with an early onset after transplantation.⁷ Non-destructive PTLTDs occur more commonly in younger patients with no prior EBV infection. Polymorphic PTLTDs show features of architectural effacement, but do not meet diagnostic criteria of any known lymphoid neoplasms. The histological features of monomorphic PTLTDs on the other hand will meet the diagnostic criteria of known lymphoid neoplasms as seen in immunocompetent patients. The survival outcomes and treatment modalities of PTLTDs are highly variable. The non-destructive PTLTDs generally have good clinical response to just reduction in immune suppression. On the other hand, polymorphic and monomorphic PTLTDs, besides reduction in immune suppression, may need additional treatment like radiotherapy, surgery, anti-CD20 rituximab drug therapy, immunotherapy and other forms of systemic chemotherapy for disease control. The type of treatment is dependent on the exact PTLTD histological subtype, age, disease stage and other blood investigations like lactate dehydrogenase levels.

Plasmacytoma PTLTD is a rare form of monomorphic B-cell PTLTD that is more commonly reported in adult patients. The histological features are similar to those of plasmacytoma in immunocompetent patients, and are usually composed of sheets of plasma cells with light chain restriction. Some cases also have smaller foci of large lymphoid cells with nucleoli resembling plasmablasts. Many cases of plasmacytoma PTLTD have had good clinical outcomes even with limited

therapy.^{10,11} The resolution of plasmacytoma PTLTD in our case, attributed to the patient's immune reconstitution, supported those findings. Various treatment options of plasmacytoma PTLTD include reduction of immune suppression, localised radiation, surgery and systemic chemotherapy based on myeloma protocols.^{10,11}

In summary, we present a case of EBV negative monomorphic B-cell plasmacytoma PTLTD in a patient who received AHSCT for advanced stage CHL. It is important for pathologists and treating haematologists to diagnose and recognise PTLTDs, as their first line of treatment is usually reduction in immune suppression. Some forms of PTLTD also have good prognosis even with minimal to limited therapy.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest to disclose.

Ming Liang Oon¹, Gwyneth Soon¹, Li Mei Poon², Shi Wang¹

¹Department of Pathology, National University Hospital, National University Health System, Singapore; ²Department of Haematology-Oncology, National University Hospital, National University Health System, Singapore

Contact Shi Wang.

E-mail: shi_wang@nuhs.edu.sg

1. Awaya N, Adachi A, Mori T, *et al.* Fulminant Epstein-Barr virus (EBV)-associated T-cell lymphoproliferative disorder with hemophagocytosis following autologous peripheral blood stem cell transplantation for relapsed angioimmunoblastic T-cell lymphoma. *Leuk Res* 2006; 30: 1059–62.
2. Jenkins D, DiFrancesco L, Chaudhry A, *et al.* Successful treatment of post-transplant lymphoproliferative disorder in autologous blood stem cell transplant recipients. *Bone Marrow Transpl* 2002; 30: 321–6.
3. Zambelli A, Lilleri D, Baldanti F, *et al.* Hodgkin's disease as unusual presentation of post-transplant lymphoproliferative disorder after autologous hematopoietic cell transplantation for malignant glioma. *BMC Cancer* 2005; 5: 109.
4. Lohrlich CA, Nevill TJ, Barnett MJ, *et al.* Development of a biologically distinct EBV-related lymphoproliferative disorder following autologous bone marrow transplantation for an EBV-negative post-renal allograft Burkitt's lymphoma. *Leuk Lymphoma* 2000; 39: 195–201.
5. Lones MA, Kirov I, Said JW, *et al.* Post-transplant lymphoproliferative disorder after autologous peripheral stem cell transplantation in a pediatric patient. *Bone Marrow Transplant* 2000; 26: 1021–4.
6. Peniket AJ, Perry AR, Williams CD, *et al.* A case of EBV-associated lymphoproliferative disease following high-dose therapy and CD34-purified autologous peripheral blood progenitor cell transplantation. *Bone Marrow Transplant* 1998; 22: 307–9.
7. Swerdlow SH, Campo E, Harris NL, *et al.* *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*. Revised 4th ed. Lyon: IARC Press, 2017; 453–62.
8. Nelson BP, Nalesnik MA, Bahler DW, *et al.* Epstein-Barr virus-negative post-transplant lymphoproliferative disorders: a distinct entity? *Am J Surg Pathol* 2000; 24: 375–85.
9. Heslop HE. How I treat EBV lymphoproliferation. *Blood* 2009; 114: 4002–8.
10. Trappe R, Zimmermann H, Fink S, *et al.* Plasmacytoma-like post-transplant lymphoproliferative disorder, a rare subtype of monomorphic B-cell post-transplant lymphoproliferation, is associated with a favourable outcome in localized as well as in advanced disease: a prospective analysis of 8 cases. *Haematologica* 2011; 96: 1067–71.
11. Perry AM, Aoun P, Culter DW, *et al.* Early onset, EBV(-)PTLTD in pediatric liver-small bowel transplantation recipients: a spectrum of plasma cell neoplasms with favorable prognosis. *Blood* 2013; 121: 1377–83.

DOI: <https://doi.org/10.1016/j.pathol.2019.05.008>