Anatomical Pathology

A REVIEW OF EXTERNAL QUALITY ASSURANCE PROGRAM PERFORMANCE IN NON-GYNAECOLOGICAL CYTOLOGY AND PD-L1 FOR NSCLC

Julia Pagliuso

The Royal College of Pathologists of Australasia Quality Assurance Programs (RCPAQAP), Anatomical Pathology, St Leonards, NSW, Australia

The Royal College of Pathologists of Australasia Quality Assurance Programs (RCPAQAP) are accredited against International Standard (ISO/IEC) 17043 Conformity assessment – General requirements for proficiency testing. This standard requires evaluation and commentary of participant performance including variation between participants and comparisons between previous testing rounds. To implement these requirements for the Cytopathology non-gynaecological programs in 2019, methods from the Anatomical Pathology diagnostic programs were utilised. Methods for case selection, evaluation of participants and results from the first year will be discussed.

In 2017, the RCPAQAP began collaborating with the United Kingdom National External Quality Assessment Services for Immunocytochemistry and In-situ hybridisation (UK NEQAS ICC and ISH) for PD-L1 for non small cell lung cancer (NSCLC) technical proficiency testing. In 2018 there were 22 RCPAQAP participants enrolled in this program and in 2019, enrolments increased to 40 participants. Results from 2018/19 will be discussed and comparisons with the broader UK NEQAS participants will be made.

The RCPAQAP issued a PD-L1 for NSCLC interpretation pilot and results from this pilot will also be discussed along with other updates and developments for Anatomical Pathology and Cytopathology programs in 2020.

CRYOBIOPSY VERSUS SURGICAL LUNG BIOPSY IN THE DIAGNOSIS OF INTERSTITIAL LUNG DISEASE (COLDICE STUDY): IS BIGGER ALWAYS BETTER?

Annabelle Mahar¹, Wendy A. Cooper^{1,2,3}, Jeffrey L. Myers⁴, Christopher Grainge⁵, Tamera J. Corte^{2,6}, Jonathan P. Williamson^{7,8}, Michael P. Valley^{2,8}, Paul J. Torzillo^{2,6}, Martin J. Phillips^{8,9}, Ganesh Raghu¹⁰, Edmund M. T. Lau^{2,6}, Lauren K. Troy^{2,6}, on behalf of The COLDICE Study Investigators

¹Tissue Pathology and Diagnostic Oncology, NSW Pathology, Royal Prince Alfred Hospital, Sydney, NSW, Australia; ²Sydney Medical School, University of Sydney, Sydney, Australia; ³School of Medicine, Western Sydney University, Sydney, NSW, Australia; ⁴Michigan Medicine, University of Michigan, Ann Arbor, MI, USA; ⁵John Hunter Hospital, New Lambton Heights, NSW, Australia; ⁶Royal Prince Alfred Hospital, Camperdown, NSW, Australia; ⁷Liverpool Hospital, Liverpool, NSW, Australia; ⁸Macquarie University Hospital, NSW,

Australia; ⁹Sir Charles Gairdner Hospital, Nedlands, WA, Australia; and ¹⁰University of Washington, Seattle, WA, USA

Background: Cryobiopsy is a less invasive technique than surgical lung biopsy (SLB) for sampling lung tissue for pathological diagnosis of interstitial lung disease (ILD), however, the diagnostic accuracy of cryobiopsy is unknown.

Methods: We investigated the diagnostic agreement of cryobiopsy and SLB in 65 patients with ILD who underwent both procedures sequentially, in a prospective multicentre diagnostic study. Biopsies were assessed independently by 3 pathologists and a consensus diagnosis was reached for guideline-directed patterns of definite or probable usual interstitial pneumonia (UIP), indeterminate for UIP, and alternative diagnosis, as well as specific histological diagnosis. Each biopsy was subsequently discussed at a multidisciplinary meeting (MDM) with clinical, radiological and pathological data.

Results: Diagnostic agreement between cryobiopsy and SLB was 70.8% following pathological assessment alone (weighted κ 0.70, 95% CI 0.55–0.86) and 76.9% following MDM (κ 0.62, 0.47–0.78). Cryobiopsies were concordant with SLB diagnoses in 37/39 (95%) cases with high or definite diagnostic confidence at MDM. Interobserver agreement between 3 pathologists was κ =0.53 (95% CI 0.43–0.63) for guideline-directed pattern in cryobiopsies and κ =0.64 (95% CI 0.54–0.75) in SLBs.

Conclusion: There is high diagnostic agreement between cryobiopsy and SLB for diagnosis of ILD, supporting the use of cryobiopsy in clinical practice.

RE-EVALUATION OF PT STAGING CATEGORIES FOR RENAL CELL CARCINOMA

Brett Delahunt

Wellington School of Medicine and Health Sciences, University of Otago, Wellington, New Zealand

Staging of renal cell carcinoma has evolved since the initial classification of Flocks and Kadesky in 1958. Currently most cancer registries utilise the UICC or AJCC classifications, now both in their eighth editions. These classifications assign localised tumours to pT1-2 categories, with the two subdivisions being based on tumour size. pT3 tumours show regional infiltration, with renal sinus invasion (pT3a) being most frequently seen. This study was undertaken to determine the association between extrarenal tumour spread and size in a series of well-sampled clear cell renal cell carcinoma (ccRCC). In a series of 917 cases of ccRCC, 178 were >7 cm in maximum extent. Assessment of tumours >7 cm in size showed 72 (40.4%) to have renal sinus infiltration, the tumour infiltrating perirenal fat in 7 (3.9%) cases, and both in 96 (53.9%) cases. In the remaining 3 (1.7%) cases, no extrarenal extension of the tumour was seen. These 3 cases with organ-confined ccRCC were all cystic tumours. For tumours ≤7 cm in maximum extent there was an increase in the proportion of tumours showing extrarenal spread with increasing size. These findings indicate that the defining