interpretation of a T cell receptor (TCR) panel and reinforce the importance of interpreting diagnostic tests in combination with clinical details.

References

1. Bleesing JH, Nagaraj CB, Zhang K. Autoimmune lymphoproliferative syndrome. In: Adam MP, Ardinger HH, Pagon RA, *et al.*, editors. GeneReviews. 14 Sep 2006, updated 24 Aug 2017, cited Dec 2019. Seattle: University of Washington, 1993-2019.
2. Worth A, Thrasher AJ, Gasper BH. Autoimmune lymphoproliferative syndrome: molecular basis of disease and clinical phenotype. *Br J Haematol* 2006; 133: 124–40.

DIAGNOSIS OF CHRONIC GRANULOMATOUS DISEASE (CGD) BY DIHYDRORHODAMINE (DHR) TEST: A SINGLE CENTRE EXPERIENCE

B. K. Prasobhan¹, M. F. Baharin¹, M. Hishamshah¹, I. H. Ismail², S. J. Lim³, B. S. Chua⁴, M. Chok⁵, A. M. Ripen¹

¹Allergy and Immunology Research Centre, Institute for Medical Research (IMR), Kuala Lumpur, Malaysia;

²Department of Paediatrics, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia;

³Department of Paediatrics, Hospital Melaka, Malaysia;

⁴Department of Paediatrics, Hospital Raja Permaisuri Bainun Ipoh, Malaysia;

and ⁵Department of Paediatrics, Hospital Tunku Jaafar Seremban, Malaysia

Introduction: Chronic granulomatous disease (CGD) is a heterogenous group of disorders that usually manifests in the first few years of life with recurrent pyogenic infections. Clinically highly suspicious patients subjected to the dihydrorhodamine (DHR) test would allow the quantitative determination of neutrophil oxidative burst rapidly.

Aim: To describe a one year findings of CGD screening in Malaysia.

Method: A total of 52 samples received for DHR test on appointment basis over a one year period. These whole blood samples were processed using commercial reagent kit Phagoburst (Glycotope Biotechnology, Germany) according to the manufacturer's protocol. This kit contains three stimulants, *E.coli*, PMA and fMLP. Samples were analysed in a BD FACSCanto II flowcytometer.

Results: Four cases showed impaired neutrophil oxidative burst activity and four more cases were inconclusive. The only positive male patient had a strong family history and the DHR test was also done for the mother which suggested she was a carrier. One patient was also found to have an 'inflamed granuloma' on histopathological report, which is a common complication in CGD. Methicillin sensitive *Staphylococcus aureus* (MSSA) predominates most of the culture and sensitivity isolates.

Discussion: Dihydrorhodamine (DHR) test is a rapid tool to detect both the diseased patient as well as the X-linked carrier parent. There is a good correlation between NBT and DHR with DHR test being more rapid, more sensitive and able to identify the carrier states of X-linked CGD. The carrier pattern is observed due to the heterogeneous population of neutrophils having abnormal and normal phagocytic function. All the inconclusive DHR tests were from the East Malaysian hospitals and the sample integrity was the main contributing factor. Optimal results are obtained in samples transported within 24 hours of blood collection.

References

1. Dimitrova G, Bunkall C, Lim D, *et al.* Comparison of two methods for the diagnosis of chronic granulomatous disease – neutrophil oxidative burst measured by the nitroblue tetrazolium slide test versus the dihydrorhodamine 123 flow cytometric assay. *NZ J Med Lab Sci* 2013; 67: 45–51.
2. Bortoletto P, Lyman K, Camacho A, *et al.* Chronic granulomatous disease : a large, single-center US experience. *Pediatr Infect Dis J* 2015; 34: 1110–14.

THE LYMPHOCYTE TRANSFORMATION TEST, FLOW CYTOMETRY AND HLA TYPING IN VANCOMYCIN INDUCED DRESS: A CASE SERIES FROM SOUTH AUSTRALIA

Matthew Kruppenacher¹, Khalida Perveen², Antonio Ferrante^{1,2}, Pravin Hissaria¹

¹SA Pathology, Adelaide, Australia; and ²Robinson Research Institute, University of Adelaide, Adelaide, Australia

Background: The lymphocyte transformation test (LTT) has been utilised to identify the culprit drug in drug hypersensitivity reactions in a research setting, but has not been validated as a diagnostic test. HLA-A*32:01 is a recently recognised risk factor for vancomycin induced drug reaction with eosinophilia and systemic symptoms (DRESS).

Methods: For LTT, peripheral blood mononuclear cells were incubated with vancomycin (10, 100, 1000 µg/mL). After 5 days, cells were pulsed with tritiated thymidine (³HTdR) and harvested after 6 h to determine the response. Results were expressed as stimulation index (SI), = disintegrations per minute (DPMs) in the presence of drug divided by the absence. An SI >2 was considered positive. CD4⁺ T cells were examined for intracytoplasmic IL-4 and IFNγ by flow cytometry. Phytohemagglutinin was a positive control. HLA-A*32:01 expression was also determined.

Results: Two patients showed proliferation in response to vancomycin in a dose-dependent manner (SI of 5 and 85). One expressed HLA-A*32:01, the other is pending testing. IL-4, but not IFNγ, was increased in CD4⁺ T cells. Further patients are undergoing testing (to be presented).

Conclusion: LTT and HLA-typing show promise for investigating vancomycin induced DRESS. Drug-induced elevation of IL-4 is consistent with a Th2-mediated pathophysiology for DRESS. Further work is underway to standardise this test for routine diagnostic use.

A NOVEL MASS SPECTROMETRIC-BASED PROTEOMIC APPROACH CHALLENGES THE CURRENT PARADIGM ON CRYOGLOBULIN TYPING IN THE DIAGNOSTIC LABORATORY

Adrian Y. S. Lee^{1,2,3}, Dimitra Beroukas^{1,2}, Alex Colella^{1,2}, Timothy Chataway², Anthony J. F. Smith³, Scott W. Graf⁴, Paul A. J. Russo⁵, Tom P. Gordon^{1,2,3,*}, Jing Jing Wang^{1,2,*}

¹SA Pathology (Flinders Medical Centre), Bedford Park, SA, Australia; ²College of Medicine and Public Health, Flinders University, Bedford Park, SA, Australia; ³Flinders Medical Centre, Bedford Park, SA, Australia; ⁴Wakefield House Rheumatology, Adelaide, SA, Australia; and ⁵Chandlers Hill Surgery, Happy Valley, SA, Australia; *Equal senior authors