relates to the relevant departments, in this case immunology, haematology and clinical chemistry. Test-specific information such as test names, result standardisation, reference intervals and comments belong to individual departments.

Current evidence indicates that there is wide variation in pathology reports at all of these levels. The solutions to these issues also lie in different locations and will rely on relevant professional organisation, wider pathology organisations (RCPA), manufacturers, standards organisations and, most importantly, individual laboratories.

HYPOGAMMAGLOBULINAEMIA AFTER RITUXIMAB THERAPY

Darren M. Roberts^{1,2}

¹*Department of Renal Medicine and Transplantation, St Vincent's Hospital, Darlinghurst, NSW, Australia; and*

²*Department of Clinical Pharmacology and Toxicology, St Vincent's Hospital, Darlinghurst, NSW, Australia*

Rituximab is an anti-CD20 chimeric monoclonal antibody with an expanding role in the treatment of autoimmune diseases. There are also increasing data regarding its adverse effect profile. Of particular interest is infectious complications of which hypogammaglobulinaemia is a risk factor. Hypogammaglobulinaemia was not reported in RCTs but there are mounting data regarding its association with rituximab therapy. Observational studies have identified it in up to 20% of patients receiving rituximab (usually much less) but the incidence and severity is poorly defined due to differing approaches to monitoring and diagnostic criteria. Hypogammaglobulinaemia may occur within 6 months or many years after initiation. Proposed risk factors include other immunosuppression (e.g., cyclophosphamide), pre-existing hypogammaglobulinaemia and/or the condition being treated. Clinical implications vary from none (spontaneous resolution) to recurrent infections. In a small number, immunoglobulin replacement therapy is required for persistent hypogammaglobulinaemia with recurrent severe infections. Resolution of hypogammaglobulinaemia is commonly reported, including in those in whom it is severe and persistent in the context of ongoing rituximab. More data may better define the risks and time course of hypogammaglobulinaemia but in the meantime it is prudent to monitor for its development in all patients treated with rituximab with regular quantification of immunoglobulin concentrations.

OPTIMUM LAB PRACTICES FOR DIAGNOSIS AND MONITORING OF MYELOMA

Joanna Sheldon

Protein Reference Unit, South West London Pathology, St George's Hospital, London, UK

There is a multitude of standards on the diagnosis of myeloma (and other B cell malignancies) that become relevant when a patient is referred to Clinical Haematology with a high clinical index of suspicion. However, there are no robust standards to enable laboratories to optimise their practices even though the abnormal pathology results, particularly those from the protein

section, often drive these referrals. There are many elements to consider: what samples to analyse, which tests to offer and which methods to use. The expected methodological characteristics should be defined, e.g., sensitivity of the immunoglobulin quantification or the presence of a robust antigen excess check in the serum free light chain method. The process should be appropriate for the spectrum of samples that may arrive in the protein section of the lab, e.g., a process to the collection and analysis of samples that are hyperviscous or have cryoproteins. Every lab should have an appropriate training and competence process with an evaluation stage for all staff working in this heavily interpretative area of the lab. There should be agreed reporting schemes that are consistently used. Finally, there must be a robust process for rapid communication of clinically critical results.

RECOMMENDATIONS FOR STANDARDISED REPORTING OF PROTEIN ELECTROPHORESIS

Nilika Wijeratne^{1,2}

¹*Chemical Pathology, Dorevitch Pathology, Heidelberg, Vic, Australia; and* ²*Department of Medicine, Monash University, Clayton, Vic, Australia*

Protein electrophoresis is vital in the diagnosis, monitoring and prognostication of monoclonal gammopathies. While clinical guidelines have recommendations on required testing for these monoclonal gammopathies, they do not recommend the exact methodologies that clinical laboratories should use for the quantification and reporting of monoclonal proteins. The inter-laboratory variation in quantification and reporting of paraproteins by protein electrophoresis may impact patient care if the patient uses different laboratory services for monitoring their disease response. The need for laboratory harmonisation has accelerated recently with the introduction of electronic health records in Australia.

While there is variability across all aspects of the quantification and reporting of protein electrophoresis, it is particularly apparent for certain aspects including co-migrating paraproteins, small abnormal bands and urine protein electrophoresis. Harmonisation requirements for reporting of protein electrophoresis include terminology, units, methods of quantification, reporting formats and interpretative commenting. The 2012 recommendations for standardised reporting of protein electrophoresis in Australia and New Zealand¹ and the 2019 proposed addendum² aimed to harmonise the laboratory practices related to paraprotein testing. However, the heterogeneity of measurements and reporting of paraproteins still exists.³

References

1. Tate JR, Caldwell G, Daly J, *et al.* Recommendations for standardized reporting of protein electrophoresis in Australia and New Zealand. *Ann Clin Biochem* 2012; 49: 242–56.
2. Tate JR, Smith JD, Wijeratne N, *et al.* Proposed addendum to 2012 recommendations for standardised reporting of protein electrophoresis in Australia and New Zealand. *Clin Biochem Rev* 2019; 40: 23–30.
3. Wijeratne N, Tate JR, Wienholt L, *et al.* Report of the survey conducted by RCPAQAP on current practice for paraprotein and serum free light chain measurement and reporting: a need for harmonisation. *Clin Biochem Rev* 2019; 40: 31–42.