VERRUCIFORM XANTHOMA BEHAVING BADLY

Marie Bertrand-Philippe, Ben Ryan

Sullivan Nicolaides Pathology, Bowen Hills, Qld, Australia

Verruciform xanthoma is an uncommon, asymptotic, mucocutaneous lesion which occurs predominantly in the oral mucosa of patients in their 5th to 7th decades. Other extraoral sites include skin of the anogenital area. It presents as a well circumscribed, rough, granular or pebbly plaque and can mimic a squamous cell carcinoma macroscopically. Recurrence is extremely rare. Histologically, verruciform xanthoma is characterised by a papillary epithelium with acanthosis and parakeratosis invaginating between the epidermal projection, as well as elongated rete ridges which sometimes coalesce at their base. Within the connective tissue papillae there is a collection of large xanthoma cells.

Forty-four cases of verruciform xanthoma have been reported in our laboratory over a 10 year period representing <0.001% of the dermatological cases. Although 38 of 44 (86%) of the lesions occurred in the common sites (oral and anogenital), six cases were present at atypical sites including ear, wrist, thumb and leg. We are presenting these atypical cases of verruciform xanthoma to highlight the importance of considering this diagnosis when reporting a lesion mimicking a SCC in locations where verruciform xanthomas are not usually present.

OESTROGEN RECEPTOR EXPRESSION IN A RARE CASE OF LEIOMYOMA OF DEEP SOFT TISSUE – A CASE REPORT

Ramzi Amin, Clarence Teo, Nicholas Koh

Department of Pathology, Tan Tock Seng Hospital, Singapore

Uterine leiomyoma is a common smooth muscle neoplasm. Gynaecologic type leiomyoma of deep soft tissue, on the other hand, is rare and commonly occurs in the retroperitoneum or abdominal cavity of young to middle aged women. We report a 42-year-old female who initially had hysterectomy performed for uterine leiomyoma at another institution but was also noted to have a 30 cm retroperitoneal mass. She was then referred to our institution where resection of the mass revealed a well-circumscribed tumour with histology features of a leiomyoma, supported by the expression of smooth muscle markers with immunohistochemical staining. Additionally, the lesional cells were both oestrogen-receptor and progesterone-receptor positive, a typical feature of this lesion which suggests derivation from hormonally sensitive smooth muscle cells in this extra-uterine location. Awareness of this benign entity is important to avoid a misdiagnosis of leiomyosarcoma so that unnecessary additional treatment can be avoided.

HISTOPATHOLOGY OF THE PLACENTA IN CONGENITAL SYPHILIS

Jamie Bishop, Jessica Matich

Histology Department, Middlemore Hospital, Auckland, New Zealand

Syphilis, a disease with a long and controversial past, has recently shown a resurgence in many countries. Cases of congenital syphilis are now being reported in New Zealand,

having previously been very rare prior to 2012. Congenital syphilis may lead to perinatal death or long term morbidity. The traditional histological triad seen in placentas consists of enlarged hypercellular villi, proliferative villous vasculature, and villitis. Necrotising funisitis, with a 'baberpole' appearance at macroscopy, is another common finding. *Treponema pallidum* immunohistochemistry may assist in the diagnosis. The previous rarity of the disease means that many pathologists may not be very familiar with the histopathological features. We present two recent cases of congenital syphilis, with placental histology, as a reminder. Recognition of congenital syphilis is important in aiding diagnosis, leading to appropriate follow up and treatment for both mother and infant.

References

1. Ministry of Health. *National Syphilis Action Plan: An action plan to stop the syphilis epidemic in New Zealand*. Wellington: Ministry of Health, 2019.
2. Tsimis ME, Sheffield JS. Update on syphilis and pregnancy. *Birth Defects Res* 2017; 109: 347–52.
3. Genest DR, Choi-Hong SR, Tate JE, *et al*. Diagnosis of congenital syphilis from placental examination: comparison of histopathology, Steiner stain, and polymerase chain reaction for *Treponema pallidum* DNA. *Hum Pathol* 1996; 27: 366–72.

ROSETTE-FORMING GLIONEURONAL TUMOUR IN A PATIENT WITH NEUROFIBROMATOSIS TYPE 1 – A CASE REPORT

Jessica Bonavita, Peter Robbins

Department of Anatomical Pathology, PathWest Laboratory Medicine, QEII Medical Centre, Perth, WA, Australia

Rosette-forming glioneuronal tumour (RGNT) is a rare primary CNS neoplasm originally described arising in the fourth ventricle, and displaying a distinctive biphasic appearance of neurocytic and astrocytic components. Patients with neurofibromatosis type 1 (NF1) can have a variety of CNS manifestations, including neoplastic, and RGNT has been described in four cases to date. This case study reports a patient with NF1 who underwent resection of a left cerebellar lesion. Histology showed a lesion dominated by synaptophysin positive, neurocytic cells in a myxoid background, forming perivascular pseudorosettes and true rosettes surrounding a central eosinophilic fibrillary core. A second component of Olig2+, oligodendroglial-like cells was also noted in a nondescript glial background. Due to its readily recognisable appearance, a diagnosis of RGNT was made. This is now a fifth case report of RGNT, an already rare tumour, in a patient with NF1 and raises the interesting possibility of a potential association between the two.

SOLID PAPILLARY CARCINOMA OF THE BREAST WITH REVERSE POLARITY – A CASE REPORT OF A RARE ENTITY WITH A NOVEL IDH2 MUTATION

Jessica Bonavita, Celia Green, Ramela Rajakaruna,

Fabienne Grieco-Iacopetta, Benhur Amanuel, Felicity Frost

Department of Anatomical Pathology, PathWest Laboratory Medicine, QEII Medical Centre, Perth, WA, Australia

Solid papillary carcinoma with reverse polarity (SPCRP) is a rare breast carcinoma bearing resemblance to the tall cell variant of papillary thyroid carcinoma (PTC). These tumours have been