no cystic changes or intraepithelial eosinophilic materials/calcifications. Dentinogenic ghost cell tumour is a benign but locally infiltrating neoplasm, mainly composed of ameloblastoma-like epithelium, arranged in sheets, islands, and nests within a background of mature fibrous connective tissue, and characteristically containing variable amount of ghost cells and dentinoid material.¹ Absence of ghost cells and presence of immature fibrous connective tissue in our Case 2 readily ruled out dentinogenic ghost cell tumour. Lastly, the morphology of this tumour, together with its clinical and radiological features, is highly suggestive of POT. However, compared with classic POTs, our Case 2 displayed some unusual features, including florid epithelial proliferation of ameloblastic epithelium with cystic degeneration, prominent intraepithelial eosinophilic materials/calcifications, and focal dentinoid deposits. In previously reported cases of POT, although the surface epithelium may form invaginated strands, all tumours were characterised by a predominant mesenchymal proliferation.² In our Case 2, the epithelial invagination was also present in most areas, but in areas the epithelial proliferation was so prominent that we believe this is true extension of the epithelial component. To our knowledge, the presence of dentinoid in association with POT has not been described thus far.^{1–8,10}

In our opinion, Case 2 might be reasonably identified as POT with previously undescribed morphological features. First, although the ameloblastic epithelium exhibited florid proliferation extending deep into the mesenchymal component in Case 2, it was continuous with the surface epithelium. The finding of florid epithelial proliferation may not be surprising, considering that previous studies have shown both epithelial and mesenchymal activity in POTs.² Second, although not reported in the original study, small foci of intraepithelial calcifications have been described in three cases of POT, as well as our first case.^{6–8} A more recent case reported by Poomsawat *et al.* contained many eosinophilic masses of varying degree of calcifications, similar to our Case 2.¹⁰ Third, dentinoid deposition is unspecific and has been identified in a wide variety of lesions, such as odontogenic keratocyst, glandular odontogenic cyst, adenomatoid odontogenic tumour, dentinogenic ghost cell tumour, ameloblastoma, odontogenic fibroma, odontogenic carcinoma, and so on.^{1,11,12} The nature of dentinoid is still poorly understood; it may represent hyalinisation or induction of the adjacent connective tissues as a result of signalling from the odontogenic epithelium.¹ Therefore, the presence of dentinoid should not be considered a diagnosis of exclusion of POT. The importance of its recognition is to avoid misdiagnosis with other odontogenic tumours. In fact, our Case 2 was originally diagnosed as ameloblastic fibro-odontoma in 2006, mainly due to the presence of these enameloid and dentinoid materials.

Combining our cases with those from the literature, we consider that POTs may span a broad histopathological spectrum with a varying number and differentiation of the epithelial component, ranging from a layer of cuboidal epithelial cells resembling the internal enamel epithelium on one end, to ameloblastic epithelium with florid proliferation on the other. Various amounts of amorphous eosinophilic material resembling enameloid and calcified material within the epithelial component can occur. Dentinoid, albeit rare and limited, may be present. Future case reports may shed more light on this new lesion.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest to disclose.

Ming Zeng¹, Xinming Chen^{1,2}, Xiaolong Guo^{1,3}, Shadong Yang^{1,2}

¹The State Key Laboratory Breeding Base of Basic Science of Stomatology (Hubei-MOST) and Key Laboratory of Oral Biomedicine Ministry of Education, School and Hospital of Stomatology, Wuhan University, Wuhan, Hubei, China; ²Department of Pathology, School and Hospital of Stomatology, Wuhan University, Wuhan, Hubei, China; ³Department of Oral Radiology, School and Hospital of Stomatology, Wuhan University, Wuhan, Hubei, China

Contact Dr Shadong Yang.

E-mail: ysd170811@whu.edu.cn

1. Mosqueda-Taylor A, Neville BW. Primordial odontogenic tumour. In: El-Naggar AK, Chan JKC, Grandis JR, Takata T, Slootweg PJ, editors. *WHO Classification of Tumours of the Head and Neck*. 4th ed. Lyon: IARC Press, 2017; 223–4.
2. Bologna-Molina R, Pereira-Prado V, Sánchez-Romero C, *et al.* Primordial odontogenic tumor: a systematic review. *Med Oral Patol Oral Cir Bucal* 2020; 25: e388–94.
3. Mosqueda-Taylor A, Pires FR, Aguirre-Urizar JM, *et al.* Primordial odontogenic tumour: clinicopathological analysis of six cases of a previously undescribed entity. *Histopathology* 2014; 65: 606–12.
4. Slater LJ, Eftimie LF, Herford AS. Primordial odontogenic tumor: report of a case. *J Oral Maxillofac Surg* 2016; 74: 547–51.
5. Ando T, Shrestha M, Nakamoto T, *et al.* A case of primordial odontogenic tumor: a new entity in the latest WHO classification (2017). *Pathol Int* 2017; 67: 365–9.
6. Mikami T, Ohashi Y, Bologna-Molina R, *et al.* Primordial odontogenic tumor: a case report with histopathological analyses. *Pathol Int* 2017; 67: 638–43.
7. Almazyad A, Li CC, Tapia ROC, *et al.* Primordial odontogenic tumour: report of two cases. *Histopathology* 2018; 72: 1221–7.
8. Bomfim BB, Prado R, Sampaio RK, *et al.* Primordial odontogenic tumor: report of a new case and literature review. *Head Neck Pathol* 2019; 13: 125–30.
9. Mikami T, Bologna-Molina R, Mosqueda-Taylor A, *et al.* Pathogenesis of primordial odontogenic tumour based on tumorigenesis and odontogenesis. *Oral Dis* 2018; 24: 1226–34.
10. Poomsawat S, Ngamsom S, Nonpassopon N. Primordial odontogenic tumor with prominent calcifications: a rare case report. *J Clin Exp Dent* 2019; 11: e952–6.
11. Mosqueda-Taylor A, Neville BW, Tatemoto Y, *et al.* Odontogenic carcinoma with dentinoid: a new odontogenic carcinoma. *Head Neck Pathol* 2014; 8: 421–31.
12. Ide F, Kikuchi K, Miyazaki Y, *et al.* Archegonous cystic odontoma is not necessarily primordial. *Head Neck Pathol* 2016; 10: 418–21.

DOI: <https://doi.org/10.1016/j.pathol.2020.04.007>

Malignant fibroadenoma with *CHEK2* mutation: a novel case

Sir,

A 34-year-old fit and well lady presented to her general practitioner complaining of a palpable left breast lump at the 12 o'clock region for 2 years. An ultrasound was requested and showed a 23 mm oval, hypoechoic lesion with minor vascularity. The differential diagnoses included a fibroadenoma and a phyllodes tumour.

Fig. 1 H&E. (A) Benign fibroadenoma on the right transitioning to DCIS (arrow) on the left. (B) Widespread low grade DCIS. (C) DCIS and the invasive component (arrow).

She subsequently proceeded to have an ultrasound guided core biopsy which showed a benign fibroepithelial lesion consistent with a fibroadenoma. However, there were features of focal ductal hyperplasia with intracytoplasmic mucin and squamoid differentiation. These features were considered unusual and an excisional biopsy was suggested in view of the findings.

She was referred to a specialist breast surgeon for follow up. Further clinical history revealed a maternal great grandmother with breast cancer. The patient opted for an excisional biopsy. The histopathology of the excision showed a complex fibroadenoma with areas of low grade ductal carcinoma *in situ* (DCIS) (Fig. 1A,B) and a focal area of early G2 invasive ductal carcinoma (IDC) (Fig. 1C) of non-special type confirmed with immunohistochemistry (Fig. 2). Hormonal receptors were ER positive, PR positive and HER2 negative. Given that there was early IDC minimally clear of the resection margins, the patient proceeded to have a sentinel lymph node biopsy and re-excision, both of which show no evidence of malignancy. Interestingly, despite the fibroadenoma showing extensive epithelial proliferation with *in situ* carcinoma and invasive carcinoma, the adjacent breast parenchyma was non-proliferative.

The case was discussed at the multidisciplinary meeting, which recommended tamoxifen, radiotherapy and genetic testing due to her young age. The genetic testing panel included *BRCA1*, *BRCA2*, *TP53*, *STK11*, *PTEN*, *CHEK2*, *PALB2* and *ATM* genes. The result identified a germline mutation in checkpoint kinase 2 (*CHEK2*) 1100delC. The patient subsequently opted for bilateral prophylactic mastectomies which were negative for malignancy.

Fibroadenoma is one of the most common benign breast tumours. The reported incidence of carcinoma arising within a fibroadenoma is rare and ranges from 0.0125% to 0.3% with a peak age range of 42–44 years old which is approximately 20 years later than the peak age of a usual fibroadenoma.¹ Malignancies occurring in fibroadenomas are usually carcinoma *in situ* and very rarely invasive carcinomas.²

It is postulated that carcinoma involving a fibroadenoma may arise from the adjacent breast tissue and infiltrate the fibroadenoma or arise within the fibroadenoma as well as the adjacent breast tissue. The presence of fibroadenoma or even complex fibroadenoma in itself is not considered an independent risk factor for breast cancer.³ Only a small proportion of fibroadenomas are ever sampled or excised, with the majority of cases followed up conservatively.

Of the small quantity of cases sampled, the reported rates of preoperative malignancies diagnosed (via cytology or core

biopsy) are very rare.^{4,5} This is due to the heterogeneity of the lesion and extensive sampling is necessary because the areas of interest can be very focal.

Studies have demonstrated that benign breast diseases (BBD) in fibroadenomas do not lead to an increased risk of malignancy when compared to BBD in general breast tissue.³ To date no known histological features on core biopsies have been shown to be associated with increased risk of malignancy in fibroadenomas.

Reported clinico-radiological risk factors warranting an excision biopsy in fibroadenomas include symptomatic lesions, newly appearing or enlarging mass, immobile or poorly circumscribed mass, presence of microcalcifications, age >35 years, size >2.5 cm and atypical biopsy features.^{6,7} Our case showed BBD on the core biopsy and there was family history of breast cancer two generations prior. Nevertheless, no studies have shown an association between family history of breast cancer and fibroadenoma with malignancy.

The rarity of the lesion precludes systematic assessment and hence there are no management guidelines established. Most authors agree that fibroadenomas with carcinoma behave like breast cancer of the same stage and should be managed similarly. However, the prognosis is reported to be more favourable than conventional types of breast carcinoma, which raises the question on the need for radiation therapy in completely excised lesions.⁸

In recent times, molecular studies have assisted in the understanding of familial breast cancer arising in non-

Fig. 2 Immunohistochemistry staining with CK5/6 showing loss of myoepithelial cells.

BRCA1/BRCA2 patients. One of the genes that have been implicated is the *CHEK2* gene. The significance of changes in the *CHEK2* gene is not well understood. Based on current information, it is indicated that the gene is a risk modifier for developing breast cancer.

Studies have shown that individuals with a *CHEK2* gene fault have increased risk (20–25%) for developing breast cancer in general and cases with a previous diagnosis have an even greater risk (30–40%) to represent with further breast cancer in their lifetime.^{9,10} This increase in cancer risk is also dependent on personal and family history. Other studies have suggested that *CHEK2* carriers also show an increased risk of cancer in the colon, prostate, thyroid and ovary.¹¹

CHEK2 is a tumour suppressor gene that encodes for serine/threonine kinase and plays a major role in the regulation of the cell cycle, apoptosis and DNA repair. Early studies suggest *CHEK2* may have prognostic implications, showing that the use of chemotherapy in combination with *CHEK2* inhibitors has beneficial effects by sensitising tumour cells to the drugs.¹²

One study has shown that there is significantly increased frequency of germline mutations in DCIS under the age of 40. These findings are consistent with our patient's demographics, and similarly in line with the reported peak age of fibroadenoma with malignancy in the literature.

None of the cases of fibroadenoma with malignancy reported in the literature mention molecular findings. It is unknown how many of these cases are associated with germline mutations. Further studies are required to assess this subgroup of patients.

To the authors' knowledge, we report the first case of *CHEK2* mutation present in a fibroadenoma with *in situ* and invasive malignancy. This case illustrates the importance of molecular testing in unusual breast cancers. The identification of germline mutations such as *CHEK2* has current and future merit in management and prognostic implications involving the patient and family members.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest to disclose.

Daniel Ching¹, Shadi Pirasteh¹, Bhavya Vora², Richard Martin³, Jespall Gill¹

¹Western Diagnostics Pathology, Myaree, Perth, WA, Australia; ²King Edward Memorial Hospital, Subiaco, Perth, WA, Australia; ³Mr Richard Martin General Surgery Subiaco, Subiaco, Perth, WA, Australia

Contact Dr Daniel Ching.

E-mail: Danielching.lh@gmail.com

1. Pick PW, Iossifides IA. Occurrence of breast carcinoma within a fibroadenoma. A Review. *Arch Pathol Lab Med* 1984; 108: 590–6.
2. Greenberg R, Skornick Y, Kaplan O. Management of breast fibroadenomas. *J Gen Intern Med* 1998; 13: 640–5.
3. Nassar A, Visscher DW, Degnim AC, Frank RD, Vierkant RA, Frost M. Complex fibroadenoma and breast cancer risk: a mayo clinic benign breast disease cohort study. *Breast Cancer Res Treat* 2015; 153: 397–405.
4. Psarianos T, Kench JG, Ung OA, Bilous AM. Breast carcinoma in a fibroadenoma: diagnosis by fine needle aspiration cytology. *Pathology* 1998; 30: 419–21.
5. Yano Y, Ueno E, Kamma H, et al. Non-invasive lobular carcinoma within a fibroadenoma, a preoperative diagnosed case. *Breast Cancer* 2001; 8: 70–3.

6. Hubbard JL, Cagle K, Davis JW, Kaups KL, Kodama M. Criteria for excision of suspected fibroadenomas of the breast. *Am J Surg* 2015; 209: 297–301.
7. Baker KS, Monsees BS, Diaz NM, Destouet JM, McDivitt RW. Carcinoma within fibroadenomas: mammographic features. *Radiology* 1990; 176: 371–4.
8. Abe H, Hanasawa K, Naitoh H, Endo Y, Tani T, Kushima R. Invasive ductal carcinoma within a fibroadenoma of the breast. *Int J Clin Oncol* 2004; 9: 334–8.
9. Schmidt MK, Hogervorst F, van Hien R, et al. Age- and tumor subtype-specific breast cancer risk estimates for *CHEK2**1100delC carriers. *J Clin Oncol* 2016; 34: 2750–60.
10. Cybulski C, Wokolorczyk D, Jakubowska A, et al. Risk of breast cancer in women with a *CHEK2* mutation with and without a family history of breast cancer. *J Clin Oncol* 2011; 29: 3747–52.
11. Apostolou P, Papatouliou I. Current perspectives on *CHEK2* mutations in breast cancer. *Breast Cancer* 2017; 9: 331–5.
12. Cybulski C, Lubiński J, Wokolorczyk D, et al. Mutations predisposing to breast cancer in 12 candidate genes in breast cancer patients from Poland. *Clin Genet* 2015; 88: 366–70.

DOI: <https://doi.org/10.1016/j.pathol.2020.04.009>

Primary breast telangiectatic osteosarcoma: a rare entity and a potential diagnostic pitfall

Sir,

Telangiectatic osteosarcoma (TOS) is a rare osteosarcoma subtype with an incidence of 2–12% among all cases of osteosarcoma.¹ Considering the rarity, reliable data documenting mammary TOS are scarce. So far, there are only two references which report that breast TOS arises as a result of the very unusual phenomenon of further malignant differentiation within a pre-existing benign or malignant neoplasm of the breast.^{2,3} In this article, we describe a rare case showing the unequivocal morphology of TOS of breast without breast precursor lesions. To the best of our knowledge, this case represents the first case of TOS in breast without breast associated diseases.

A 60-year-old woman presented to our centre for evaluation of a painful lump in her left breast for the previous 10 days. Her medical history was significant for hypertension and chronic bronchitis. Physical examination revealed a fixed mass, 4×3 cm, in the upper outer quadrant of her left breast. Ultrasonography (US) showed a hypoechoic, cystic mass with relatively regular luminal outline and numerous fluid levels at 3 o'clock, 2.5 cm from the nipple. Biopsy revealed a high-grade malignant tumour.

Lumpectomy was performed. Macroscopically, the surgical specimen measured 4.5×4.0×4.0 cm (Fig. 1A). Sectioning revealed a haemorrhagic cystic mass with blood clots (Fig. 1B). Microscopically (Fig. 2), spaces were blood-filled and separated by septa containing highly malignant mesenchymal cells with osteoid. Bone calcification was absent. Mitoses were obvious. Immunohistochemical staining was performed using antibodies against cytokeratins (AE1/AE3, Cam5.2, CK7, CK5/6, CK14, CK17, 34βE12 and CK20), epithelial membrane antigen (EMA), P53, EA, CD31, CD34, ERG, S100, SOX10, Melan A, CD10, CD117, calponin, desmin, CD68, CD163, CD99, IDH1, SATB2, oestrogen receptor (ER), progesterone receptor (PR) and human epidermal growth factor receptor 2 (HER2).