that involvement at diagnosis was confined to the abdominal cavity with involvement of the retroperitoneum extending into the root of the mesentery, and no evidence of intrathoracic disease.

Histopathologically, KLA is distinguished by foci of 'kaposiform' spindled lymphatic endothelial cells, often arranged in parallel fashion either as dispersed, poorly marginated clusters or anastomosing strands.³ Further immunohistochemical staining, such as with PROX-1, D2-40, ERG, CD31 or CD34 can confirm the presence of these lymphatic endothelial cells.^{7,9,10} This is distinct from kaposiform haemangioendothelioma (KHE), another disorder associated with abnormal lymphatic channels and spindled cells, however the histological architecture is of more defined solid confluent tumour, often with microthrombi.³ Clinically, KHE differs further as the lesions are typically unifocal vascular tumours presenting in infancy, which may involve skin and soft tissues, retroperitoneum or viscera,³ with larger lesions commonly associated with thrombocytopenia resulting in consumptive coagulopathy (Kasabach–Merritt phenomenon).

There is currently no standard treatment for KLA and management is directed at symptom control and management of complications such as drainage of pericardial or pleural effusions.^{3,7,9} Pharmacotherapy options such as sirolimus, an mTOR inhibitor, have been utilised in other vascular anomalies with promising results.¹¹ Croteau *et al.* reported clinical improvement in three patients following sirolimus therapy,³ and other case studies have also reported reduction in lesion size or improvement in symptoms^{6,8,9} or stable disease.⁷ One patient in a case series by Fernandes *et al.*, showed initial response to sirolimus and a 5 month period of remission before subsequent reaccumulation of pleural and pericardial effusion, and development of epidural haemorrhage resulting in fatal cardiac arrest.⁷ Limited survival and prognostic data exist for KLA; however, Croteau *et al.* reported a 5 year survival of 51% and an overall survival of 34%.³ Mean interval between diagnosis and death was reported as 2.75 years with cardiorespiratory failure being the most common cause.³ Of note, there was one patient in the series who presented as an adult, never required any interventions, and denied any issues on follow-up.³

In conclusion, KLA is a rare but emerging entity classically presenting as intrathoracic lesions in young patients. However, this diagnosis should also be considered in patients with infiltrating masses in the retroperitoneum even when intrathoracic disease is not present, as demonstrated with this case.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest to disclose.

George Chen¹, Nicole S. Graf², Tony Pang³

¹Department of Surgery, Westmead Hospital, Westmead, NSW, Australia; ²The Children's Hospital Westmead, Westmead, NSW, Australia; ³The University of Sydney, Sydney, NSW, Australia

Contact Dr George Chen.

E-mail: gzuo_chen@hotmail.com

1. Debelenko L, Marler J, Perez-Atayde AR, *et al.* Kaposiform lymphangiomatosis (KLA): an aggressive variant of lymphangiomatosis. *Lab Invest* 2004; 84: 269.

2. Perez-Atayde AR, Croteau SE, Trenor CC, *et al.* Kaposiform lymphangiomatosis: a new entity distinct from generalized lymphatic anomaly, kaposiform hemangioendothelioma, lymphedema-associated angiosarcoma, and lymphangioma-like kaposi sarcoma. International Society of Vascular Anomalies Meeting, 2012, Malmo, Sweden.
3. Croteau SE, Kozakewich HP, Perez-Atayde AR, *et al.* Kaposiform lymphangiomatosis: a distinct aggressive lymphatic anomaly. *J Pediatr* 2014; 164: 383–8.
4. Safi F, Gupta A, Adams D, *et al.* Kaposiform lymphangiomatosis, a newly characterized vascular anomaly presenting with hemoptysis in an adult woman. *Ann Am Thorac Soc* 2014; 11: 92–5.
5. Goyal P, Alomari AI, Kozakewich HP, *et al.* Imaging features of kaposiform lymphangiomatosis. *Pediatr Radiol* 2016; 46: 1282–90.
6. Raam MS, Festekjian A, Elkhunovich MA. Point-of-care thoracic ultrasonography in the diagnosis and management of kaposiform lymphangiomatosis. *Pediatr Emerg Care* 2016; 32: 888–91.
7. Fernandes VM, Fargo JH, Saini S, *et al.* Kaposiform lymphangiomatosis: unifying features of a heterogeneous disorder. *Pediatr Blood Cancer* 2015; 62: 901–4.
8. Radzikowska E, Blasinska-Przerwa K, Szolkowska M, *et al.* Kaposiform lymphangiomatosis with human papillomavirus infection. *Am J Respir Crit Care Med* 2017; 195: e47–8.
9. Wang Z, Li K, Yao W, *et al.* Successful treatment of kaposiform lymphangiomatosis with sirolimus. *Pediatr Blood Cancer* 2015; 62: 1291–3.
10. Ordonez NG. Immunohistochemical endothelial markers: a review. *Adv Anat Pathol* 2012; 19: 281–95.
11. Triana P, Dore M, Cerezo VN, *et al.* Sirolimus in the treatment of vascular anomalies. *Eur J Pediatr Surg* 2017; 27: 86–90.

DOI: <https://doi.org/10.1016/j.pathol.2020.07.014>

Non-mammary implant associated anaplastic large cell lymphoma

Sir,

Anaplastic large cell lymphoma (ALCL) constitutes less than 3% of all non-Hodgkin lymphomas (NHL).¹ Implant associated ALCL was first reported by Keech and Creech² in 1997 in a patient with a breast implant. In the last two decades, several case reports and small case series of breast implant associated ALCL (BIA-ALCL) have been reported in the literature and this is now a provisional entity in the current World Health Organization (WHO 2017) classification of hematolymphoid neoplasms.³ The currently estimated incidence of BIA-ALCL in the United States is 33 per one million persons with textured breast implants.⁴ The textured implants are associated with a higher risk of disease development as compared to saline implants.⁵ In 2010, Palraj *et al.*⁶ reported ALCL in association with a stainless steel implant for complex right tibia fracture, and recently Mendes *et al.*⁷ documented ALCL following augmentation gluteoplasty by silicone implant. In the literature, only two cases of non-mammary implant associated extranodal ALCL have been reported. We present a case of a male with a history of oral cavity squamous cell carcinoma for which he underwent marginal mandibulectomy and device implantation (titanium; smooth surface) and developed ALCL 4 years later. To the best of our knowledge, this is the second case of non-silicone prostheses associated ALCL and the third case of implant associated ALCL at the non-mammary site.

A 63-year-old man, with well controlled diabetes and hypertension, had a non-healing ulcer over the left buccal mucosa in 2016. On oral examination an ulcero-proliferative growth measuring 3 × 3 cm was seen in the left buccal mucosa involving the retromolar trigone posteriorly and reaching the

gingiva buccal sulcus (cT3N0M0). No cervical lymphadenopathy was noted. The mucosal biopsy revealed squamous cell carcinoma, well differentiated (Fig. 1A). Contrast enhanced computed tomography (CT) scan showed the involvement of buccinator muscle by growth, however there was no evidence of erosion of the underlying mandible or maxilla. The patient underwent wide local excision of the left buccal mucosa with marginal mandibulectomy, upper alveolectomy, left submandibular gland excision, and left modified neck dissection type III. This was followed by free artery-based forearm flap and split thickness skin graft (STSG) for secondary defects. The next day of surgery, a six-hole titanium plate and a six-screw implant was performed as a result of surgical site clot formation (Fig. 1B and C). On pathology examination of the specimen, the depth of invasion was 1 cm (pT2NoMx) and all resected margins and the cervical lymph nodes were free of tumour. He received 30 cycles (60 Gy) of local radiotherapy over the course of one month. Three years later, he developed jaw pain and his metallic implant became exposed in the oral cavity. Magnetic resonance imaging (MRI) was performed with suspicion of recurrence or osteonecrosis and revealed an ill-defined soft tissue mass with mild contrast enhancement at the post-operative site. On further follow-up, the exposed implant was removed by surgery. At this time, the surgeon felt a nodule measuring 1.5×1.0 cm in the upper part of the left cheek along with skin mobilization (Fig. 1D). Excision biopsy of the nodule was carried out.

The wide local excision specimen was skin covered and measured $2.0 \times 1.5 \times 1.0$ cm. The attached skin was relatively unremarkable. The cut surface revealed an ill-defined grey-white lesion measuring $1.5 \times 1.0 \times 0.6$ cm that was firm in consistency and located within the dermis and subcutis but not invading the epidermis. Microscopically, the lesion was

composed of predominantly large atypical cells in a dyscohesive growth pattern forming vague nodules in the dermis and subcutis. Tumour cells had large irregular nuclei with prominent nucleoli. The characteristic hallmark cells were also present showing kidney-shaped nuclei and brisk mitosis (Fig. 2A). The background showed mild fibrosis, and chronic inflammation. On immunohistochemistry, leukocyte common antigen (LCA) showed membranous positivity. CD3 and CD20 immunostains were negative but the background reactive T-cells were highlighted by CD3 (Fig. 2C). CD30 showed diffuse strong and uniform immunoreactivity in the large cells in the form of membranous and golgi complex, however ALK immunostain was negative (Fig. 2B,D). MUM1 depicted strong, diffuse, nuclear immunostaining (Fig. 2E). CD38, CD138, S100p, HMB45, pan-cytokeratin and synaptophysin, were all negative. In addition, p63 immunostain showed diffuse nuclear positivity (>90%) in these large cells (Fig. 2F). *In situ* hybridization for Epstein–Barr virus was negative. Fluorescence *in situ* hybridization (FISH) did not reveal *DUSP22* rearrangement (Fig. 2G). On histopathology, the case was diagnosed as ALCL, ALK-negative with TP63 expression. The probable cause could be non-silicon (titanium) implant associated. The patient has undergone three cycles of CHOP chemotherapy (cyclophosphamide, doxorubicin, vincristine, and prednisolone) to date and interim disease assessment is pending.

There are only a few cases of lymphoma reported in association with metallic prostheses or osteosynthetic materials, consisting mainly of large B-cell lymphomas and only one case of large T-cell lymphoma, i.e., ALCL.^{6,8} The exact pathogenesis behind this association is not yet understood, especially those occurring at non-mammary sites. Presently, orthopaedic implants are typically composed of titanium or cobalt-chromium alloys, stainless steel, titanium, or tantalum,

Fig. 1 (A) Squamous cell carcinoma, well differentiated, of left buccal mucosa (H&E). (B) Left lateral plain radiographic view of mandible showing six hole titanium plate and six screw implant. (C) On removal, the smooth surfaced titanium implant. (D) A 1.5×1.0 cm well demarcated, firm, nodule seen at surgical site of implant.

Fig. 2 Non-mammary implant associated anaplastic large cell lymphoma (NMIA-ALCL). (A) Characteristic morphology of ALCL, common pattern. The tumour cells are large with high N:C ratio. Occasional hallmark cells are present (H&E). The tumour cells are (B) diffusely positive for CD30; (C,D) negative for CD20, and ALK; (E,F) diffusely positive for MUM1 and p63 immunostains. (G) Fluorescence *in situ* hybridization did not reveal *DUSP22* rearrangement.

with or without a polyethylene articulating surface.⁹ The index case had undergone titanium implant placement in the mandibular region following surgery for oral squamous cell carcinoma.

Non-mammary implant associated ALCL was not documented within the literature until Palraj *et al.*⁶ reported the first case in 2010. They described a case of primary extranodal ALCL developing in the setting of joint prosthesis where the immunocompetent elderly patient underwent implantation of stainless steel plate for open reduction internal fixation (ORIF) of a complex tibia fracture and developed ALCL 7 years later at the site of implant. Kellogg *et al.*⁹ looked retrospectively through the literature and gathered data on association of non-Hodgkin lymphoma or ALCL with mammary and non-mammary (heart valve prostheses, blood vessel prostheses, orthopedic fixation devices, arthroplasty, artificial pacemakers, and indwelling catheters) implants. Their focus was on non-mammary implants showing association with non-Hodgkin lymphoma. On thorough search they found only one case of non-mammary implant associated ALCL which was reported by Palraj *et al.* as mentioned above. Mendes *et al.*⁷ described another non-mammary implant associated ALCL in a 63-year-old female after augmentation gluteoplasty with silicone implants. Both the above reported ALK-negative ALCL were morphologically and immunophenotypically similar to BIA-ALCL. The only discordance was with the type of implant used: stainless steel in the joint prosthesis case and silicone implant in the gluteoplasty.

All BIA-ALCL patients showed a 'triple negative' genetic profile—negative for *ALK*, *DUSP22* and *TP63*

rearrangements—and were characterised by mutational and gene expression profiles consistent with aberrant activation of the JAK/STAT pathway.³ The present case was negative for ALK immunoexpression and *DUSP22* rearrangement but showed diffuse p63 immunoreactivity. The latter positivity on immunohistochemistry is considered a screening test which needs further confirmation by FISH for *TP63* rearrangement (not done in our case). The previous two reported cases were not investigated for *DUSP22* or *TP63* rearrangements.

The pathogenesis for mammary or non-mammary prostheses associated ALCL is yet unknown and needs to be explored. There are many hypotheses considered like chronic inflammation as the potential etiological agent in view of the ability of implants to promote T-cell infiltration and immune response.¹⁰ Other possibilities considered include high expression of interleukin-2 and interleukin-6 receptors; strong activation of signal transducer and activator of transcription 3 (STAT3) signaling, likely causing autocrine production of IL-6; chronic bacterial antigen stimulation and sustained T-cell production; and high levels of IL-13 and IgE as part of the ALCL microenvironment causing chronic allergic reaction.¹¹ Palraj *et al.*⁶ suggested chronic inflammatory reaction to a particular antigenic stimulus (the implant) that evolved to lymphoma through clonal transformation and/or genetic mutation of a particular subset of lymphocytes, while Mendes *et al.*⁷ considered the role of implant surface in ALCL genesis (textured are more prone to lymphoma genesis than smooth). The role of chronic inflammation rather than implant surface can be considered more in our case.

In implant associated lymphomas, prosthesis explantation and complete excision of any residual mass are the recommended approaches. Systemic chemotherapy is generally recommended for patients with locally advanced and disseminated diseases, mainly using treatment protocols routinely adopted for systemic ALCLs (e.g., CHOP, CHOEP, etc.). Limited data are available regarding the treatment of non-mammary implant associated ALCL. The first documented joint prosthesis associated ALCL patient was lost to follow-up while no adjuvant chemotherapy was considered in gluteal implant associated ALCL, however the patient was disease free. Our patient has received chemotherapy (CHOP regime) and was doing well at last follow-up.

In conclusion, oncologists and pathologists should be more observant in patients who undergo silicone or non-silicone device implantation as these surgical procedures will continue to rise in various non-neoplastic and neoplastic lesions. Though the morphology and immunophenotyping of mammary and non-mammary implant associated ALCL is similar to ALK-negative ALCL, such cases need to be further investigated for *DUSP22* or *TP63* rearrangements as the latter carries dismal prognosis. The exact biological relationship between the implant device and ALCL needs to be addressed and explored at the molecular level.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest to disclose.

Mayur Parkhi¹, Charanpreet Singh², Pankaj Malhotra², Amanjit Bal¹

¹Department of Histopathology, Post Graduate Institute of Medical Education and Research (PGIMER), Chandigarh, India; ²Department of Internal Medicine and Clinical Hematology, Post Graduate Institute of Medical Education and Research (PGIMER), Chandigarh, India

Contact: Dr Amanjit Bal.
E-mail: docaman5@hotmail.com

- Kim B, Roth C, Chung KC, *et al.* Anaplastic large cell lymphoma and breast implants: a systematic review. *Plast Reconstr Surg* 2011; 127: 2141–50.
- Keech Jr JA, Creech BJ. Anaplastic T-cell lymphoma in proximity to a saline-filled breast implant. *Plast Reconstr Surg* 1997; 100: 554–5.
- Swerdlow SH, Campo E, Harris NL, *et al.*, editors. *WHO classification of tumours of haematopoietic and lymphoid tissues*. Revised 4th edition. Lyon: IARC Press, 2017.
- Doren EL, Miranda RN, Selber JC, *et al.* Epidemiology of breast implant-associated anaplastic large cell lymphoma. *Plast Reconstr Surg* 2017; 139: 1042–50.
- Collett DJ, Rakhorst H, Lennox P, *et al.* Current risk estimates of breast implant-associated anaplastic large cell lymphoma in textured breast implants. *Plast Reconstr Surg* 2019; 143: 30S–40S.
- Palraj B, Paturi A, Stone RG, *et al.* Soft tissue anaplastic large T-cell lymphoma associated with a metallic orthopedic implant: case report and review of the current literature. *J Foot Ankle Surg* 2010; 49: 561–4.
- Mendes Jr J, Mendes Maykeh VA, Frascino LF, *et al.* Gluteal implant-associated anaplastic large cell lymphoma. *Plast Reconstr Surg* 2019; 144: 610–3.
- Sanchez-Gonzalez B, Garcia M, Montserrat F, *et al.* Diffuse large B-cell lymphoma associated with chronic inflammation in metallic implant. *J Clin Oncol* 2013; 31: e148–51.
- Kellogg BC, Hiro ME, Payne WG. Implant-associated anaplastic large cell lymphoma: beyond breast prostheses. *Ann Plast Surg* 2014; 73: 461–4.
- Hu H, Jacombs A, Vickery K, *et al.* Chronic biofilm infection in breast implants is associated with an increased T-cell lymphocytic infiltrate:

implications for breast implant-associated lymphoma. *Plast Reconstr Surg* 2015; 135: 319–29.

- Marra A, Viale G, Pileri SA, *et al.* Breast implant-associated anaplastic large cell lymphoma: a comprehensive review. *Cancer Treat Rev* 2020; 84: 101963.

DOI: <https://doi.org/10.1016/j.pathol.2020.06.022>

Metachronous and clonally related classic Hodgkin lymphoma and nodular lymphocyte predominant Hodgkin lymphoma

Sir,

Classic Hodgkin lymphoma (CHL) and nodular lymphocyte predominant Hodgkin lymphoma (NLPHL) are two distinct subtypes of Hodgkin lymphoma by the World Health Organization (WHO) classification.¹ They share some common clinical and pathological features, but are different in immunophenotype, inflammatory background, and genetic characteristics. The coexistence of CHL and NLPHL in the same patient is exceedingly rare.² We present a case of CHL where complete remission was achieved after treatment, but clonally related NLPHL developed subsequently.

A previously healthy 14-year-old male presented with generalised lymphadenopathy. He had no B symptoms. His uncle had a history of CHL. The computed tomography (CT) scan revealed lymphadenopathy in the left neck, bilateral supraclavicular, left axilla, anterior mediastinum and right hilum, and multiple low attenuating lesions in the spleen. The positron emission tomography (PET) scan confirmed glucose avid lymphadenopathy of these areas with a maximum SUV of 16.1 at right hilum. An excisional biopsy of left supraclavicular lymph node showed a nodular infiltrate dissected by thick fibrotic bands (Fig. 1A). The infiltrate was composed of scattered large Hodgkin/Reed–Sternberg (HRS) cells in a background of small lymphocytes, eosinophils, histiocytes, and plasma cells (Fig. 1B). The HRS cells were positive for PAX5 (weak), CD15 and CD30, and negative for CD20, CD45 and EMA (Fig. 1C–F). *In situ* hybridisation for EBV encoded small RNA (EBER) was negative. The findings were consistent with nodular sclerosis CHL. The cytogenetic analysis revealed a complex karyotype with an abnormal hypertriploid clone characterised by additional unknown material on Xq, 1q, 2p, and 7q, isochromosomes of 7p and 10q, gain of 1, 2, 10, 19, and 5–6 marker chromosomes and loss of 6 and 17 {78–79,XY,add(X)(q?24),+1,add(1)(q32)x2,+2,add(2)(p16)x2,-6,+add(7)(q31)x2,i(7)(p10),+10,i(10)(q10)x2,-17,+19,+5–6mar[5]/46,XY[10]}. The patient's bone marrow was not involved by the CHL. He was designated stage IIIS CHL and was treated per Children's Oncology Group (COG) protocol AHOD0031 for intermediate risk Hodgkin disease. He received four cycles of ABVE-PC (adriamycin, bleomycin, vincristine sulfate, etoposide, prednisone and cyclophosphamide). Post-chemotherapy the CT scan showed resolution of lymphadenopathy and splenic mass. The PET scan showed no convincing evidence of residual lymphoma. He subsequently received radiation therapy of 2100 cGy to the neck, mediastinum and spleen. He was in clinical remission.