Table 1 Approach to identifying children with FH, modified from European Atherosclerosis Society child and adolescent guidelines⁶

LDL cholesterol	Action required
≥5.0 mmol/L	Repeat lipid profile with TSH, liver and kidney function tests and urine protein to exclude common secondary causes of elevated LDL-c. Refer to regional lipid service if repeat LDL-c ≥5.0 mmol/L.
4.0–4.9 mmol/L	Determine if a parent has a history of elevated LDL-c or premature CVD (males <55 years, females <60 years). Repeat lipid profile with TSH, liver and kidney function tests and urine protein to exclude common secondary causes of elevated LDL-c. Refer to regional lipid service if repeat LDL-c ≥4.0 mmol/L and a parent has elevated LDL-c or premature CVD.
3.5–3.9 mmol/L	Repeat lipid profile with TSH, liver and kidney function tests and urine protein to exclude common secondary causes of elevated LDL-c. Enquire if a parent has been diagnosed with FH. Refer to regional lipid service if repeat LDL-c ≥3.5 mmol/L and a parent has been diagnosed with FH.

CVD, cardiovascular disease; FH, familial hypercholesterolaemia; LDL-c, low density lipoprotein cholesterol; TSH, thyroid stimulating hormone.

guidelines⁶ is presented in Table 1. Approximately 10% of children would require additional questions regarding family history and/or repeat lipid testing, which should not overly burden a practising dermatologist.

In conclusion, dermatologists can play a significant role in the detection of FH by incorporating opportunistic screening for FH into their guidelines for managing isotretinoin therapy. However, further research is required to explore this potential method of opportunistic FH detection.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest to disclose.

Aaron J. Frederiks¹, John R. Burnett^{1,2,3}, Gerald F. Watts^{1,2}, Sujith P. Kumarasinghe¹, Damon A. Bell^{1,2,3,4}

¹School of Medicine, University of Western Australia, Perth, WA, Australia; ²Lipid Disorders Clinic, Cardiometabolic Service, Department of Cardiology, Royal Perth Hospital, Perth, WA, Australia; ³Department of Clinical Biochemistry, PathWest Laboratory Medicine WA, Royal Perth Hospital and Fiona Stanley Hospital Network, Perth, WA, Australia; ⁴Department of Clinical Biochemistry, Clinipath, Osborne Park, WA, Australia

Contact A/Prof Damon A. Bell.
E-mail: damon.bell@health.wa.gov.au

1. Nordestgaard BG, Chapman MJ, Humphries SE, *et al.* Familial hypercholesterolaemia is underdiagnosed and undertreated in the general population: guidance for clinicians to prevent coronary heart disease: consensus Statement of the European Atherosclerosis Society. *Eur Heart J* 2013; 34: 3478–90.

2. Lan NSR, Martin AC, Brett T, *et al.* Improving the detection of familial hypercholesterolaemia. *Pathology* 2019; 51: 213–21.
3. Sjouke B, Kusters DM, Kindt I, *et al.* Homozygous autosomal dominant hypercholesterolaemia in The Netherlands: prevalence, genotype-phenotype relationship, and clinical outcome. *Eur Heart J* 2015; 36: 560–5.
4. Luirink IK, Wiegman A, Kusters DM, *et al.* 20-year follow-up of statins in children with familial hypercholesterolaemia. *N Engl J Med* 2019; 381: 1547–56.
5. Watts GF, Gidding S, Wierzbicki AS, *et al.* Integrated guidance on the care of familial hypercholesterolaemia from the International FH Foundation. *Int J Cardiol* 2014; 171: 309–25.
6. Wiegman A, Gidding SS, Watts GF, *et al.* Familial hypercholesterolaemia in children and adolescents: gaining decades of life by optimizing detection and treatment. *Eur Heart J* 2015; 36: 2425–37.
7. Bell DA, Hooper AJ, Bender R, *et al.* Opportunistic screening for familial hypercholesterolaemia via a community laboratory. *Ann Clin Biochem* 2012; 49: 534–7.
8. The Australasian College of Dermatologists. *The Australasian College of Dermatologists Position Statement: Isotretinoin for Treatment of Acne*. Sydney: The Australasian College of Dermatologists; 2018. <https://www.dermcoll.edu.au/wp-content/uploads/ACD-Position-Statement-isotretinoin-June-2018.pdf>
9. Barth JH, Macdonald-Hull SP, Mark J, *et al.* Isotretinoin therapy for acne vulgaris: a re-evaluation of the need for measurements of plasma lipids and liver function tests. *Br J Dermatol* 1993; 129: 704–7.
10. Rodondi N, Darioli R, Ramelet AA, *et al.* High risk for hyperlipidemia and the metabolic syndrome after an episode of hypertriglyceridemia during 13-cis retinoic acid therapy for acne: a pharmacogenetic study. *Ann Intern Med* 2002; 136: 582–9.
11. Bender R, Edwards G, McMahon J, *et al.* Interpretative comments specifically suggesting specialist referral increase the detection of familial hypercholesterolaemia. *Pathology* 2016; 48: 463–6.

DOI: <https://doi.org/10.1016/j.pathol.2020.07.007>

Rapid detection of enteric pathogens in a medical assistance team by real-time multiplex PCR

Sir,

Diarrhoeal diseases acquired through contact with unhygienic conditions and contaminated food pose an ongoing risk to the health of travellers around the world.¹ A range of pathogens are known to cause diarrhoeal disease in travellers with *Escherichia coli* pathotypes and *Shigella* considered some of the most important.^{1–3} ‘Traveller’s diarrhoea’ is usually associated with travel from developed to underdeveloped areas of the world for a period of <2 weeks.¹ While the effects of diarrhoeal disease are typically mild (e.g., tolerable, no impact on activities), moderate (substantial symptoms; impacts on activity) to severe (dysentery; incapacitating) symptoms may develop.¹ Military and non-military personnel deployed to underdeveloped regions from developed countries face similar problems to the average traveller² with the cumulative time lost having the potential to significantly impact operations.⁴ In deployed settings, accurate and rapid surveillance and early treatment are thus essential to minimise loss of duty days.⁵

In December 2019, an Australian medical assistance team (AUSMAT) was internationally deployed to a South Pacific island nation, to assist local health services in managing an ongoing outbreak of viral illness amongst the population. The 32 person team consisted of civilian doctors, nurses, paramedics and logistics specialists with expertise in a range of skills including infectious diseases and microbiology. Mission diagnostic equipment included a BioFire FilmArray

System (BioMérieux, USA) transported in a Pelican Case (Pelican, USA) as personal luggage on deployment, to provide additional rapid diagnostic testing capabilities if required. All medical assistance team staff resided in shared accommodation (two per room) with each room having an ensuite facility. All staff dined together at the hotel common dining area. Over the course of the 4-week period, seven deployed staff presented for medical review with severe water diarrhoea without the presence of blood. One individual reported faecal incontinence. Affected individuals also reported the presence of nausea, fatigue and malaise. Individuals were typically severely ill for at least 24 hours and then less well for a period of 72 hours. Most affected individuals were treated empirically with azithromycin after the onset of illness. Prior to the completion of testing, affected individuals suspected that diarrhoeal disease was acquired from contaminated foods sourced in the hotel restaurant. A number of the medical team were also regularly dealing with severely unwell patients prior to their diarrhoeal disease symptoms. However, it should be noted that at least two individuals with symptoms had no contact with patients.

Stool samples from three individuals were made available for diagnostic testing using the BioFire FilmArray Gastrointestinal GI Panel (BioMérieux), which provides simultaneous rapid nucleic acid detection of 22 gastrointestinal pathogens, including bacteria, viruses and several eukaryotic parasites, in 1 hour. This multiplex polymerase chain reaction (PCR) panel has previously been shown to be highly sensitive and specific for the range of pathogens tested.⁶

Testing was established within the existing hard stand laboratory facility and specimens were handled in a class 2 safety cabinet.

The clinical notes and diagnostic test results for the three patients tested are summarised in Table 1. All three individuals returned positive results for sapovirus with two individuals also found to be PCR positive for enteroaggregative *Escherichia coli*. On the basis of these results, azithromycin usage was ceased for all symptomatic patients. Infection control strategies were reinforced including isolating patients reporting diarrhoeal symptoms into single rooms. The sapovirus identification provided further evidence for a food source and the likely breakdown of hygienic practices in the hotel restaurant. Hotel staff were subsequently counselled on effective hand hygiene practices.

Sapoviruses are recognised agents of acute gastroenteritis, with clinical symptoms including diarrhoea, vomiting, nausea, myalgia, fever and malaise frequently reported

lasting between 1–4 days.⁷ Outbreaks have been reported in a variety of settings with transmission occurring via the faecal-oral route and typically via exposure to sapovirus-positive faeces contaminating surfaces, food and water sources.⁷

AUSMAT are typically deployed internationally for a period of 2 weeks and are expected to be self sufficient with rations normally used as part of team sustenance. More recently Australian responses have deployed to environments where reliance on locally prepared meals has been required. It is in this environment, AUSMAT experienced its first gastroenteritis outbreak. AUSMAT members consist of senior, highly specialised individuals expected to work in difficult and autonomous environments. Members are considered a specialised resource and are not easily replaced. Safety and welfare of AUSMAT members is managed aggressively and proactively from initial team member registration through to the conclusion of the deployment. A variety of tools, such as extended vaccinations, personality profile assessments, daily electronic heat surveillance, mental health and welfare screens are used along with access to point of care tests for infectious or non-communicable diseases. A member of the team is assigned as a team medical officer for confidentiality and timely support to team members and has, at their disposal, any relevant clinical equipment brought by the team to support diagnosis and treatment of the team member.

The application of rapid molecular techniques such as multiplex PCRs is emerging as an important adjunct to traditional diagnostic methods for the aetiological diagnosis of acute gastroenteritis,³ regardless of the patient cohort affected. In deployed settings, existing hard stand laboratory pathology services may be resource limited or not accessible for testing of samples from deployed staff. In this case, the ability to use this multiplex PCR system for team member assessment allowed us to implement an immediate management plan for all other cases as well as minimising further outbreaks, since the infectious period and potential risks to vulnerable populations is known for the pathogen detected. This is particularly valuable on AUSMAT deployments where there is very little human reserve available to support the operation. As such, the use of multiplex PCR as a 'force protection' tool to rapidly diagnose and provide epidemiological understanding to support the management of an outbreak of disease in the deployed medical assistance team was unexpected but equally valuable. The major limitation of the multiplex system used in this study (Biofire) was that only one sample could be tested at a time. Nevertheless, the results of this report highlight the value of this and other molecular

Table 1 Summary of patient clinical data and PCR testing results

	Case 1	Case 2	Case 3
Patient details			
Age, gender	32, Male	47, Male	38, Female
Deployment role	Logistics	Logistics	Nurse
Clinical details			
Symptoms	Severe diarrhoea	Severe diarrhoea with faecal incontinence	Severe diarrhoea with faecal incontinence
Duration	48 hours	96 hours	72 hours
Initial treatment	Azithromycin 500 mg daily; ceased after result available	Azithromycin 500 mg daily; ceased after result available	Azithromycin 500 mg daily; ceased after result available
PCR testing	Sapovirus +	Sapovirus +	Sapovirus +

diagnostic systems to protect the health and welfare of affected individuals with the secondary goal of reducing the days of lost duty. While there is a significant focus on the use of such portable systems for supporting military operations,^{2,8} it is foreseeable that such systems could be used in other settings to support medical staff in aiding deployed teams of individuals (e.g., sporting teams) affected by diarrhoeal diseases or other diseases with an infectious aetiology.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest to disclose.

James Branley^{1,2}, Abigail Trewin³, Timothy James Gray^{4,5}, Adam Polkinghorne^{1,2}

¹Department of Microbiology and Infectious Diseases, New South Wales Health Pathology, Nepean Blue Mountains Pathology Service, Penrith, NSW, Australia; ²The University of Sydney Medical School, Nepean Clinical School, Faculty of Medicine and Health, University of Sydney, Penrith, NSW, Australia; ³National Critical Care and Trauma Response Centre, Royal Darwin Hospital, Darwin, NT, Australia; ⁴The University of Sydney Medical School, Concord Clinical School, Faculty of Medicine and Health, University of Sydney, Concord, NSW, Australia; ⁵Department of Microbiology and Infectious Diseases, Concord Repatriation General Hospital, Concord, NSW, Australia

Contact Dr James Branley.

E-mail: james.branley@health.nsw.gov.au

1. Steffen R. Epidemiology of traveller's diarrhea. *J Travel Med* 2017; 24: S2–5.
2. Connor P, Porter CK, Swierczewski B, *et al.* Diarrhoea during military deployment: current concepts and future directions. *Curr Opin Infect Dis* 2012; 25: 546–54.
3. Vila J. New molecular tools in traveller's diarrhea. *J Travel Med* 2017; 24: S23–8.
4. Sanders JW, Putnam SD, Riddle MS, *et al.* Military importance of diarrhea: lessons from the Middle East. *Curr Opin Gastroenterol* 2005; 21: 9–14.
5. Riddle MS, Martin GJ, Murray CK, *et al.* Management of acute diarrheal illness during deployment: a deployment health guideline and expert panel report. *Mil Med* 2017; 182: 34–52.
6. Buss SN, Leber A, Chapin K, *et al.* Multicenter evaluation of the Biofire Filmarray gastrointestinal panel for etiologic diagnosis of infectious gastroenteritis. *J Clin Microbiol* 2015; 53: 915–25.
7. Oka T, Wang Q, Katayama K, *et al.* Comprehensive review of human sapoviruses. *Clin Microbiol Rev* 2015; 28: 32–53.
8. Lersethakarn P, Nakjarun MS, Silapong BS, *et al.* Detection of diarrhea etiology among U.S. military personnel during exercise Balikatan 2014, Philippines, using TaqMan array cards. *Mil Med* 2016; 181: e1669–74.

DOI: <https://doi.org/10.1016/j.pathol.2020.06.024>

Re: INSM1 is a novel prognostic neuroendocrine marker for luminal B breast cancer

Sir,

We read with great interest the article by Razvi *et al.* published in *Pathology*.¹ The authors evaluated insulinoma-associated protein 1 (INSM1), a crucial transcription factor with a zinc-finger motif initially isolated from a human insulinoma

phagemid cDNA library,² as a marker for neuroendocrine (NE) differentiation in infiltrating breast cancers (IBCs) in females. They demonstrated the usefulness of INSM1 expression, suggesting a favourable prognostic impact, with better disease-free survival, particularly for the luminal B phenotype, as well as having potential for stratifying NE neoplasms with different outcomes.¹ On the other hand, recently, we specifically described unusual cases with various NE phenotype mammary carcinomas in which the NE nature of the tumours was confirmed solely by INSM1, without expressions of traditional 'gold-standard' NE indicators, chromogranin (CG) A and synaptophysin (SYN).³ We also emphasised that the frequency of detecting NE differentiation in systemic neoplasms is anticipated to increase and that our observations might ultimately contribute to the development of novel treatments including molecular-targeted therapies for these frequently aggressive tumour entities.³

Herein, we ask, in the academic research conducted by Razvi *et al.*, what were the actual number and proportion of IBCs showing INSM1+/CG–/SYN– and whether cases with these immuno-profiles consistently had NE morphologies, i.e., solid and/or trabecular growth pattern, a peripheral palisading arrangement, polygonal, spindle or plasmacytoid cell appearance, finely granular, eosinophilic cytoplasm, 'salt-and-pepper' nuclear chromatin, and a highly vascular stroma, as in our reported cases.^{3,4} In fact, in the histological figure presented by the authors, an IBC with distinct INSM1 expression characteristically shows a nested formation of polygonal or focally fusiform cancer cells, with relatively abundant, eosinophilic to clear, fine-granular cytoplasm and a finely granular chromatin pattern, although the expression status of CG and SYN in this case is not shown.¹ Accordingly, the nature and significance of INSM1 as a representative diagnostic marker as well as an innovative research tool warrant greater emphasis.

In addition, Razvi *et al.* clarified INSM1+, CG+ and/or SYN+ IBCs to be relevant to luminal subtypes, showing both ER ($p < 0.001$ for all) and PR ($p \leq 0.015$) positive status, while being negative for HER2 ($p \leq 0.044$) and high molecular weight cytokeratins (CKs) ($p \leq 0.047$), whereas CD56 (neural cell adhesion molecule, NCAM) reactive cases showed contrasting clinicopathological features.¹ The only positive correlation was actually with CK14 ($p < 0.001$), and there were no relationships with ER, PR, HER2, CK5/6, or survival in luminal cancers.¹ It is well known that NCAM is generally used as an additional or supplementary NE indicator, except in specific circumstances such as the diagnosis of small cell pulmonary carcinoma.⁵ We recently demonstrated that in the breast oncology field, NCAM is a considerably less sensitive and less specific NE marker than previously assumed, because its expression is commonly found in the epithelial cells comprising normal mammary ducts and lobules which are essentially devoid of NE cells.⁶ We would also like to know the morphological findings of IBCs showing CD56 (NCAM) reactivity, especially those expressing this antibody exclusively, if possible, along with discussion of the implications and the potential utility.

Conflicts of interest and sources of funding: This work was supported by Grants-in-Aid for Scientific Research (No. 16K08654) from the Japanese Ministry of Education, Culture, Sports, Science and Technology and the National Hospital Organization (NHO) Grant (H29-NHO-01). The authors state that there are no conflicts of interest to disclose.