thought that the compounding effect of local immunosuppression and physical barrier in an enclosed environment allows the EBV infected B cells to evade immune-surveillance and proliferate in an 'immune privileged' environment (due to absence of EBV specific cytotoxic T cells.⁵ The HIF-1A expression supports the hypoxic state induced by the walled off space.⁷

The morphology and immunophenotype of the lesional cells in FA-DLBCL or CI-DLBCL can overlap with other EBV associated B cell lymphomas or lymphoproliferative disorders listed in the WHO classification, such as plasmablastic lymphoma, EBV+ DLBCL NOS, EBV-associated mucocutaneous ulcers, lymphomatoid granulomatosis and post-transplant lymphoproliferative disorders (PTLD).¹ All these entities have in some ways arisen from reduced immune surveillance for EBV, either from iatrogenic immunosuppression (e.g., medications, post-transplant setting), aging and immune senescence, or primary or hereditary immune deficiency.

The occurrence of this current lymphoma adjacent to a breast implant, raises the important pitfall for potential misclassification as breast implant-associated anaplastic large cell lymphoma (BI-ALCL). BI-ALCL is a new provisional entity in the revised 2017 WHO classification of lymphoid malignancies of T cell lineage, arising in association with a breast implant, usually as an effusion confined to the surrounding fibrous capsule or less often as a mass or axillary lymphadenopathy.¹ The pathogenesis of BI-ALCL is not well delineated with several plausible associations including chronic inflammation, textured implants and bacterial biofilm infection.¹⁰ A recent gene expression profiling study demonstrated a hypoxia signature suggesting this lymphoma arises in a unique microenvironment.¹¹ Mutations affecting the JAK-STAT signalling pathway and alterations in epigenetic modifiers have been identified in BI-ALCL.¹² Of note, our current case occurred in association with a macrotextured implant, which is more likely associated with the development of BI-ALCL. In our case, the initial H&E impression may have led to a misdiagnosis of BI-ALCL if only a limited panel of immunohistochemistry was performed or this diagnosis was not considered. BI-ALCLs are of T cell lineage as opposed to B cell origin in FA-DLBCL. Both tumours (BI-ALCL and FA-DLBCL) are MUM1 and CD30 positive. The anaplastic large cells of BI-ALCL usually show loss of pan-T-cell markers, and tumour cells of the FA-DLBCL showing a post-germinal centre phenotype may lose CD20 expression.

Here we report a unique case of EBV+ FA-DLBCL arising as an incidental finding in a breast implant capsulectomy specimen, expanding the geographical sites in which such cases have been reported. The similarities in its clinical presentation to BI-ALCL highlights the potential misclassification of this entity if it is not considered diagnostically. The cells of FA-DLBCL morphologically mimic other aggressive EBV-driven B cell lymphomas but the clinical presentation and host factors aid in the distinction, highlighting the importance of integrating the clinical and radiological/staging features in arriving at this diagnosis. As highlighted in the literature to date, this entity is more akin to a lymphoproliferation than a lymphoma.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest to disclose.

Christine Khoo¹, Christine McTigue², David J. Hunter-Smith³, Patricia Walker⁴

¹Department of Pathology, Peter MacCallum Cancer Centre, Melbourne, Vic, Australia; ²Dorevitch Pathology, Melbourne, Vic, Australia; ³Department of Plastic and Reconstructive Surgery, Peninsula Health, Melbourne, Vic, Australia; ⁴Department of Haematology, Peninsula Health, Peninsula Private Hospital, and Alfred Health, Melbourne, Vic, Australia

Contact Dr Christine Khoo.

E-mail: Christine.khoo@petermac.org

1. Swerdlow SH, Campo E, Harris NL, et al., editors. *WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues*. Revised 4th ed. Lyon: IARC, 2017.
2. Aozasa K, Takakuwa T, Nakatsuka S. Pyothorax-associated lymphoma, a lymphoma developing in chronic inflammation. *Adv Anat Pathol* 2005; 12: 324–31.
3. Lymphoma P, Cheuk W, Chan ACL, et al. Metallic implant-associated lymphoma. *Am J Surg Pathol* 2005; 29: 832–6.
4. Dojcinov S, Fend F, Quintanilla-Martinez L. EBV-positive lymphoproliferations of B- T- and NK-cell derivation in non-immunocompromised hosts. *Pathogens* 2018; 7: 28.
5. Boyer DF, McKelvie PA, De Leval L, et al. Fibrin-associated EBV-positive large b-cell lymphoma. *Am J Surg Pathol* 2017; 41: 299–312.
6. Gruver AM, Huba MA, Dogan A, Hsi ED. Fibrin-associated large B-cell lymphoma: Part of the spectrum of cardiac lymphomas. *Am J Surg Pathol* 2012; 36: 1527–37.
7. Boroumand N, Ly TL, Sonstein J, Medeiros LJ. Microscopic diffuse large B-cell lymphoma (DLBCL) occurring in pseudocysts. *Am J Surg Pathol* 2012; 36: 1074–80.
8. Kanno H, Naka N, Yasunaga Y, et al. Production of the immunosuppressive cytokine interleukin-10 by Epstein-Barr-virus-expressing pyothorax-associated lymphoma: possible role in the development of overt lymphoma in immunocompetent hosts. *Am J Pathol* 1997; 150: 349–57.
9. Chen BJ, Chapuy B, Ouyang J, et al. PD-L1 expression is characteristic of a subset of aggressive B-cell lymphomas and virus-associated malignancies. *Clin Cancer Res* 2013; 19: 3462–73.
10. Quesada AE, Medeiros JL, Clemens MW, Ferrufino-Schmidt MC, Pina-Oviedo S, Miranda RN. Breast implant-associated anaplastic large cell lymphoma: a review. *Mod Pathol* 2019; 32: 166–88.
11. Oishi N, Hundal T, Phillips JL, et al. Molecular profiling reveals a hypoxia signature in breast implant-associated anaplastic large cell lymphoma. *Haematologica* 2020; May 15: haematol.2019.245860.
12. Laurent C, Nicolae A, Laurent C, et al. Gene alterations in epigenetic modifiers and JAK-STAT signaling are frequent in breast implant-associated ALCL. *Blood* 2020; 135: 360–70.

DOI: <https://doi.org/10.1016/j.pathol.2020.09.022>

Pembrolizumab associated sarcoid-like lymphadenopathy mimicking progressive disease in Hodgkin lymphoma

Sir,

In recent years, the use of programmed cell death-1 (PD-1) checkpoint inhibitors such as pembrolizumab and nivolumab has emerged as a treatment modality in patients with various malignancies such as melanoma, non-small cell lung cancer, squamous cell carcinoma of the head and neck, urothelial carcinoma, microsatellite instability or mismatch repair deficient metastatic colorectal cancer and relapsed/refractory Hodgkin lymphoma (HL).¹ In HL, these treatments work via

abrogating the over-expression of surface PDL-1 by Hodgkin Reed–Sternberg (HRS) cells, thus reducing immunotolerance towards HRS cells and increasing HRS cell turnover.²

The common immunotherapy related adverse events (irAEs) documented with PD-1 inhibitor therapy in HL patients include rash, hypothyroidism, diarrhoea and pneumonitis.^{1,2} However sarcoid-like granulomatous reactions, including lymphadenopathy, have also been documented.^{3,4} Here, we report a 25-year-old male with relapsed/refractory HL, who developed granulomatous/sarcoid-like lymphadenopathy whilst undergoing pembrolizumab salvage therapy, with no previous history of autoimmune disease or sarcoidosis.

A 25-year-old male presented with symptoms due to a large mediastinal mass in December 2017. An urgent ultrasound-guided core biopsy was performed and the initial histopathology report was B-cell non-Hodgkin lymphoma. On this basis R-EPOCH (rituximab, etoposide, prednisolone, vincristine, cyclophosphamide, doxorubicin) was initiated due to clinical emergency. However, he developed worsening of symptoms and an enlarging mediastinal mass on fluorodeoxyglucose positron emission tomography (FDG-PET) scan 3 weeks after the first cycle of chemotherapy (Fig. 1A); hence, thoracoscopic biopsy was performed. These samples along with the initial histopathology slides were sent for second opinion to an expert lymphoma pathologist and the pathological diagnosis was revised as classical HL.

Since there was rapid disease progression after R-EPOCH (the chemotherapeutic agents constituting this regime are known to reduce disease burden in HL),⁵ salvage chemotherapy with one cycle of ICE (ifosfamide, carboplatin and etoposide) and two cycles of BrESHAP (brentuximab vendotin, etoposide, methylprednisolone, cytarabine and cisplatin) was administered, following which complete metabolic response (Deauville 1) on FDG-PET was achieved. Autologous stem cell transplantation was then performed

under carmustine, etoposide, cytarabine and melphalan (BEAM) conditioning followed by consolidation with involved field radiotherapy (36 Gy).

Six months after autologous stem cell transplantation (November 2018), new FDG-PET avid nodal lesions were found in cervical, Waldeyer's ring and mediastinum (Fig. 1B), for which thoracoscopic biopsy of subcarinal lymph nodes was performed. This confirmed recurrence of HL and pembrolizumab salvage therapy was implemented. After four cycles of pembrolizumab, progress PET-CT scan demonstrated clearance of the pre-treatment FDG-PET uptakes. However, new FDG-PET avid lymphadenopathy was demonstrated in the left supraclavicular region, left hilum, and the periportal region (Fig. 1C).

Given the new findings on FDG-PET raising the possibility of progression of HL whilst the patient was receiving pembrolizumab, our patient was referred for a clinical trial with a novel targeted agent, anti-lymphocyte activation gene 3 antibody (anti-LAG-3). As part of the clinical trial screening protocol, a biopsy of the left supraclavicular lymph node was performed. Surprisingly, the histopathology revealed multiple non-caseating granulomas resembling sarcoidosis (Fig. 2A) and no evidence of HL, with absence of CD30 positive or PAX5 positive HRS cells. CD163 positive epithelioid histiocytes were noted within the granulomas (Fig. 2B). Special stains for fungal and mycobacterial infections such as periodic acid-Schiff, silver methenamine and Ziehl–Neelsen were negative. Serum ACE levels were normal. Bronchioalveolar lavage showed benign cytology, with no increase in lymphocytes, neutrophils, eosinophils or evidence of alveolar haemorrhage.

Based on these findings, the diagnosis of sarcoid-like granulomatous lymphadenopathy as an irAE secondary to PD-1 inhibitor therapy was entertained. Prednisolone was commenced for management of the underlying inflammatory process and pembrolizumab therapy was restarted. Complete

Fig. 1 Full body fluorodeoxyglucose-positron emission tomography (FDG-PET) scans of patient at various time points of treatment. (A) FDG-PET avid bulky mediastinal mass at diagnosis. (B) New FDG-PET avid lymph node uptakes in cervical, Waldeyer's ring and mediastinum, 6 months after autograft. (C) Increased FDG-PET uptakes within the left supraclavicular lymph node, left hilum and periportal regions after four cycles of pembrolizumab salvage therapy. (D) Disappearance of all the previously increased FDG-PET uptakes after six cycles of pembrolizumab salvage therapy.

Fig. 2 Left supraclavicular lymph node microscopy. (A) Left supraclavicular lymph node showing non-caseating granulomas, H&E stain. (B) CD163 positive epithelioid histiocytes were noted within the granulomas.

metabolic response (Deauville 1) was achieved after two additional cycles of therapy (after cycle 6), with disappearance of previous areas of increased FDG-PET uptake (Fig. 1D). Later our patient underwent matched unrelated allogeneic stem cell transplantation and remained in complete metabolic remission on serial FDG-PET scans at the recent clinical review in July 2020.

The association of sarcoidosis and sarcoid-like granulomatous reactions with HL has been described in a few case reports and case series.^{6–8} In those studies, most of the patients underwent treatment with ABVD (doxorubicin, bleomycin, vinblastine and dacarbazine) or other bleomycin containing chemotherapy regimens. Exposure to bleomycin chemotherapy and also paraneoplastic syndrome have been implicated in sarcoid-like granulomatous reactions in Hodgkin lymphoma; however, the precise correlation and the pathogenetic mechanisms are unclear.^{6,7} Systemic flare of underlying sarcoidosis due to pembrolizumab treatment for Hodgkin lymphoma has been described in a case report.⁴

Non-necrotising sarcoid-like granulomas as irAEs have been documented in lymph nodes, bone marrow, lung, liver, spleen, heart, soft tissues and muscle in patients with melanoma, whilst on PD-1 inhibitor therapy.^{3,9} The irAEs related to PD-1 inhibitors are suspected to be T-cell mediated, secondary to a hyperactive immune response from PD-1 signalling inhibition.³ This may be consistent with findings of increased numbers of circulating CD4+ T-cells, CD8+ T-cells and natural killer cells in peripheral blood immunophenotyping after pembrolizumab treatment.¹⁰ It has been postulated that lysed tumour cells present neoantigens that promote a CD4+ (Th1 and Th17 cell) mediated response, with evidence from gene expression studies showing activation of interferon-gamma.^{3,10,11} With regards to development of non-necrotising sarcoid-like granulomas following PD-1 inhibitors, the interaction of monocyte derived histiocytes and CD4+ Th1 cells is postulated as a pathogenetic mechanism.³ In sarcoidosis, the production of TNF- α by sarcoid

macrophages and CD4+ T-cells is important in the formation and maintenance of sarcoidosis granulomata.¹² However, in PD-1 inhibitor induced granulomatous/sarcoid-like lesions, T-cell proliferation and PD-1 signalling inhibition may be the inciting and/or sustaining event for granulomata formation, which is unique.³ Steroid treatment has been noted to cause clearance of these lesions, with a few reports suggesting that it is possible to continue pembrolizumab therapy with steroid cover to treat the granulomatous/sarcoid-like lesions and the underlying malignancy concurrently.^{3,4}

In our patient, based on the interim FDG-PET scan after pembrolizumab, there was a strong suspicion of progression of HL. However sarcoid-like granulomatous reaction was demonstrated on the left supraclavicular lymph node biopsy four cycles (3 months) after initiation of pembrolizumab treatment and was not documented on the lymph node biopsies prior to commencement of pembrolizumab. After exclusion of other causes of granulomatous reactions including bleomycin (ABVD not administered), tuberculosis and fungal infection, pembrolizumab was deemed to be the causative factor by temporal association. Hence, we decided to offer the benefit of continuing pembrolizumab, while being aware of the possibility of progressive HL elsewhere among the new FDG-PET positive lesions in the lymph node sites in the thorax and abdomen. Further continuation of pembrolizumab with steroid cover led to complete metabolic response, which indicated that all the multifocal FDG-PET avid lymph node sites after cycle 4 of pembrolizumab (Fig. 1C) were granulomatous inflammatory lesions, rather than HL.

This case illustrates the development of multifocal nodal granulomatous sarcoid-like lesions during pembrolizumab therapy for refractory/relapsed HL, which mimicked failure of PD1 inhibitor treatment of HL on FDG-PET scans. In such situations repeat biopsies should be performed on FDG-PET avid lesions to differentiate benign sarcoid-like granulomatous lymphadenopathy from refractory/relapsed HL, the former auguring a good prognosis.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest to disclose.

Bruce Tsz Chun Chau^{1,2}, Penelope McKelvie^{3,4}, Ali Bazargan⁵, Muhajir Mohamed^{6,7}

¹Faculty of Medicine, University of Queensland, Herston, Qld, Australia; ²Department of Anatomical Pathology, Launceston General Hospital, Launceston, Tas, Australia; ³Faculty of Medicine, Dentistry and Health Sciences, The University of Melbourne, Melbourne, Vic, Australia; ⁴Department of Anatomical Pathology, St Vincent's Hospital Melbourne, Melbourne, Vic, Australia; ⁵Department of Haematology, St Vincent's Hospital Melbourne, Melbourne, Vic, Australia; ⁶School of Medicine, University of Tasmania, Hobart, Tas, Australia; ⁷Department of Haematology, Northern Cancer Service, Launceston General Hospital, Launceston, Tas, Australia

Contact Dr Muhajir Mohamed.

E-mails: muhajir.mohamed@ths.tas.gov.au, chau_bruce@hotmail.com

- Haanen J, Carbone F, Robert C, *et al.* Management of toxicities from immunotherapy: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2017; 28: iv119–42.
- Armand P, Chen YB, Redd RA, *et al.* PD-1 blockade with pembrolizumab for classical Hodgkin lymphoma after autologous stem cell transplantation. *Blood* 2019; 134: 22–9.
- Tetzlaff T, Nelson K, Diab A. Granulomatous/sarcoid-like lesions associated with checkpoint inhibitors: a marker of therapy response in a subset of melanoma patients. *J Immunother Cancer* 2018; 6: 14.
- Cotliar J, Querfeld C, Boswell W, *et al.* Pembrolizumab-associated sarcoidosis. *JAAD Case Rep* 2016; 2: 290–3.
- Hoppe R, Advani R. Hodgkin lymphoma. NCCN clinical practice guidelines in oncology. *J Natl Compr Canc Netw* 2020; 18: 755–81.
- Wirk B. Sarcoid reactions after chemotherapy for Hodgkin's lymphoma. *Clin Med Insights Case Rep* 2010; 3: 21–5.
- Cherk M, Pham A, Haydon A. 18F-fluorodeoxyglucose positron emission tomography-positive sarcoidosis after chemoradiotherapy of Hodgkin's disease: a case report. *J Med Case Rep* 2011; 5: 247.
- London J, Grados A, Ferme C, *et al.* Sarcoidosis occurring after lymphoma. *Medicine* 2014; 93: 1–13.
- Chorti E, Kanaki T, Zimmer L, *et al.* Drug-induced sarcoidosis-like reaction in adjuvant immunotherapy: increased rate and mimicker of metastasis. *Eur J Cancer* 2020; 131: 18–26.
- Maly J, Alinari L. Pembrolizumab in classical Hodgkin's lymphoma. *Eur J Haematol* 2016; 97: 219–27.
- Hiraki T, Hatanaka M, Arimura A, *et al.* Granulomatous/sarcoid-like reactions in the setting of programmed cell death-1 inhibition: a potential mimic of disease recurrence. *J Cutan Pathol* 2020; 47: 154–60.
- Cinetto F, Agostini C. Advances in understanding the immunopathology of sarcoidosis and implications on therapy. *Expert Rev Clin Immunol* 2016; 12: 973–88.

DOI: <https://doi.org/10.1016/j.pathol.2020.08.017>

Triphenotypic acute leukaemia: the contribution of multimodality laboratory diagnostic techniques

Sir,

Mixed phenotype acute leukaemia (MPAL) represents approximately 4% of all acute leukaemias, and most commonly manifests with myeloid and either B- or T-lymphoid defining markers.^{1–3} Mixed phenotype acute

leukaemias co-expressing B- and T-lymphoid markers are much less common, and those expressing a combination of B-, T- and myeloid markers ('triphenotypic acute leukaemia') are even rarer, accounting for approximately 2% MPAL, and <0.1% of acute leukaemia cases.^{1,2} Here we describe how the use of multimodality laboratory diagnostic techniques was required to confirm the diagnosis of trilineage mixed phenotype acute leukaemia, and its subsequent clinical course.

A 47-year-old man presented with dyspnoea and was found to have large pleural and pericardial effusions on chest X-ray and echocardiogram. Full blood examination revealed Hb 129 g/L, WCC $14.6 \times 10^9/L$, ANC $5.1 \times 10^9/L$, Plt $338 \times 10^9/L$ and blood film demonstrated 57% blasts with agranular cytoplasm and no Auer rods.

Bone marrow aspirate demonstrated 88% blasts (Fig. 1A) which were pleomorphic, intermediate to large in size, with high N:C ratio, scant basophilic cytoplasm, fine chromatin with occasional prominent nucleoli.

An initial acute leukaemia orientation flow cytometry panel (CD45, CD34, MPO, cytCD3, sCD3, CD7, CD19, cytCD79a) was performed to assign lineage, which identified an expanded population of cells within the progenitor gate, accounting for 80% of total white blood cells (WBCs) and comprising two distinct subpopulations which differed by their myeloperoxidase (MPO) expression. To further characterise the blast populations, our standard B-acute lymphocytic leukaemia (B-ALL), T-ALL and acute myeloid leukaemia (AML) panels were performed, in addition to a hybrid single tube 10 colour panel (CD45, CD19, CD38, CD7, CD22, CD24, CD117, CD66b, CD34, CD20) to confirm cross-lineage antigen expression. The dominant blast population (representing 90% of blasts) co-expressed B- and T-lymphoid markers (CD34+, cytCD3+, smCD3+, CD7+, CD19+dim, CD22+dim, CD79a+dim, CD24+dim, CD10–, MPO–, CD13+, CD33+), whereas the smaller population (10% of blasts) demonstrated co-expression of B- and T-lymphoid and myeloid markers (CD34+, cytCD3+, smCD3+, CD7+partial, CD19+dim, CD22+dim, CD24+dim, CD79a–, CD10–, MPO+, CD117+, CD13+br, CD33+br) (Fig. 1B).

Given the rarity of true triphenotypic leukaemia, cytochemistry was performed to confirm MPO expression which was positive in 15% of blasts (Fig. 1D). Immunohistochemistry on trephine confirmed that CD19, CD3, CD79a were positive in the majority of blasts. Immunohistochemistry for MPO and lysozyme was positive in 10% (Fig. 1C).

Conventional karyotyping demonstrated an abnormal karyotype (46,Y,add(X)(p11),del(3)(p21),add(10)(p11),-13,add(16)(q21),-18,add(22)(q13),+2mar[cp14]/46,XY[15]), with no *BCR-ABL1* fusion or *KMT2A*-rearrangement detected by fluorescence *in situ* hybridisation (FISH) and no *BCR-ABL1* transcripts detected by reverse transcription polymerase chain reaction (RT-PCR).

A next generation sequencing panel identified an *EZH2* mutation [NM_004456.4:c.1717A>G; p.(Thr573Ala)] at 74% variant allele frequency as the only mutation of the 76 genes assessed, including no evidence of *TP53* mutations. RNA sequencing was performed to further investigate gene fusions and identified a *DDX3X-MLLT10* in-frame fusion involving *DDX3X* exon 3 (NM_001356.3, located at Xp11) and *MLLT10* exon 17 (NM_004641.3, located at 10p12).

The patient commenced on Hyper-CVAD chemotherapy (cyclophosphamide, vincristine, doxorubicin, dexamethasone,