Method: 778 images of breast carcinoma slides from Sunshine Coast University Hospital were split into training, validation and test sets and labelled by a trained non-pathologist.

A ScaledYOLOv4 computer vision architecture was trained on 659 images, and the model was evaluated on the validation set (68 images), unseen by the model during training.

Results: The model demonstrated a sensitivity of 0.92 and a positive predictive value (PPV) of 0.64.

Discussion: The model's main weakness is its frequent misidentification of pyknotic cells. The model is most effective in tumours with high mitotic counts, as the ratio of mitotic figures to pyknotic cells is high.

Next steps include creating open-source databases for improved image quantity, and standardisation of data to encourage further, higher quality open-source projects.

Reference

1. Pantanowitz L, Hartman D, Qi Y, *et al.* Accuracy and efficiency of an artificial intelligence tool when counting breast mitoses. *Diagn Pathol* 2020; 15: 80.

OSSEOUS METAPLASIA OF A THYROID NODULE

Lucia Cochrane-Davis, Afaq Khan

Anatomical Pathology, Royal Darwin Hospital, Darwin, NT, Australia

While thyroid nodules commonly show calcification secondary to haemorrhage and necrosis, it is extremely rare for thyroid tissue to show true bony metaplasia. Ten cases of bony thyroid metaplasia have been reported in English literature to date, and all cases were female. Also, reports of 32 cases of thyroid extramedullary haematopoiesis (EMH) noted that ten included bony areas. All patients with bony metaplasia in EMH were female, as were all but two of the cases of thyroid EMH without bone.

We present the case of an obese 51-year-old male with no relevant medical history, who required hemithyroidectomy for a TIRADS-5 thyroid nodule which could not be biopsied. It was found to be a colloid nodule with areas of lamellar bone and fatty marrow, without extramedullary haematopoiesis. To our knowledge, this is the first reported case of osseous thyroid metaplasia in a male.

CARCINOSARCOMA OF THE UTERUS IN ASSOCIATION WITH TAMOXIFEN THERAPY – A CASE REPORT

Nimmy Cyriac¹, Iman Alash², Deepali Kamra¹

¹Dorevitch Pathology, Albury, NSW, Australia; and ²Dorevitch Pathology, Heidelberg, Vic, Australia

Tamoxifen is a nonsteroidal triphenylethyl compound which is widely used in the treatment of breast cancer. In addition to having anti-oestrogenic properties, tamoxifen also exhibits weak oestrogenic effects and acts on the uterine endometrium. Treatment with this drug may result in a spectrum of proliferative endometrial lesions, which include simple and atypical hyperplasia, hyperplasia with polyp formation (single and multiple), adenocarcinoma and rarely sarcomas. Patients on tamoxifen should be closely monitored for endometrial lesions. We present the case of a 54-year-old female with uterine carcinosarcoma arising in the setting of tamoxifen therapy.

Reference

1. McCluggage WG, McManus DT, Lioe TF, *et al.* Uterine carcinosarcoma in association with tamoxifen therapy. *BJOG* 2005; 104: 748–50.

IGG4-RELATED LIVER DISEASE, PRESENTING AS A LIVER TUMOUR (A CASE REPORT AND REVIEW OF LITERATURE)

Sara Diab, Siamak Sakhaie

Department of Anatomical Pathology, Australian Clinical Labs, Melbourne, Vic, Australia

IgG4-related disease is a multisystemic fibroinflammatory condition, although the liver involvement by the disease is rare. The disease is characterised by distinctive histopathologic features, most notably dense lymphoplasmacytic infiltrate, storiform fibrosis, and obliterative phlebitis. The diagnosis should be made only by an interdisciplinary approach to correlate clinical history, pathological picture and radiological findings. We report a case of an 80-year-old male patient presenting with a left lower lobe liver mass. An intrahepatic cholangiocarcinoma was suspected on imaging studies. The patient underwent left hemi-hepatectomy. Macroscopically there was a poorly circumscribed firm tumour measuring up to 47 mm, present within liver parenchyma. Microscopic examination showed dense hyalinised fibrotic hypocellular tissue with fibroblastic proliferation admixed with lymphoplasmacytic infiltrate with focal formation of lymphoid follicles. The entire lesion was then submitted and no malignancy was detected. IgG and IgG4 immunohistochemical stains demonstrated an increase in number of IgG4 positive cells (more than 30 per high power field), as well as an increase in the ratio of IgG4 to IgG positive plasma cells. This immunophenotype is in keeping with IgG4-related liver disease.

PRIMARY SPLENIC FOLLICULAR LYMPHOMA

Sara Diab, Justin du Plessis

Department of Anatomical Pathology, Australian Clinical Labs, Melbourne, Vic, Australia

Follicular lymphoma is primarily a nodal disease, with the spleen most commonly secondarily involved. The most common splenic haematolymphoid neoplasms are splenic marginal zone lymphoma and hairy cell leukaemia, and primary splenic follicular lymphoma is extremely rare. We report a case of a 60-year-old male patient presenting with massive splenomegaly without lymphadenopathy, with a past history of lung squamous cell carcinoma. The patient underwent splenectomy, which on gross examination revealed multiple tan granular nodules throughout the splenic parenchyma. Microscopically, there was expansion of white pulp by admixed generally small cleaved and elongate lymphocytes with scattered larger centroblast-like cells together with background small mature lymphocytes, without marginal zone formation. There were less than 5 centroblasts per high-power field. On immunohistochemistry, the atypical lymphoid cells were positive for CD79a, CD20, BCL6, BCL2 and CD10; and they were negative for CD5, CD23, and CD3. Fluorescence *in situ* hybridisation was performed, identifying the characteristic BCL2 rearrangement of follicular lymphoma, tumour graded WHO grade 1 with Ki67 proliferative index of

50%. Follicular lymphoma has potentially fatal consequences, and the patient died six months post-operatively.

PRIMARY SMALL CELL CARCINOMA OF THE BREAST

Sara Diab, Ban Sulaiman

Department of Anatomical Pathology, Gosford Hospital, NSW Health Pathology, NSW, Australia

Small cell carcinoma arising in breast is very rare. The diagnosis of primary small cell carcinoma can only be made if a non-mammary site is excluded clinically or if *in situ* component can be demonstrated histologically as in this case. We report a case of a 68-year-old female patient who presented with a left breast lump, which was diagnosed as a triple negative high grade cancer on core biopsy. The patient received neoadjuvant chemotherapy followed by a left breast lump removal (lumpectomy) with sentinel lymph node biopsy. Macroscopically, there is a well circumscribed tan to pink mass up to 24 mm. Microscopically, the breast shows nests of tumour cells. The tumour cells are small with scant cytoplasm, salt and pepper chromatin, nuclear moulding. Mitotic and apoptotic figures are plentiful. There is ductal carcinoma *in situ* (DCIS) with similar cytological features. Immunohistochemistry was performed. The tumour cells were positive for E-Cadherin, p53 and CK7, they were weakly positive for NSE and INSM1 and they were negative for CK5/6, S100 and TTF1. Ki67 score was 90%. Features are compatible with primary small cell carcinoma. Small cell carcinoma has a very aggressive behaviour. The management of the primary tumour has not been standardised, due to the rarity of this entity.

SYRINGOCYSTADENOCARCINOMA PAPILLIFERUM: A RARE ADNEXAL SKIN TUMOUR

Pranav Dorwal¹, Joel Pinczewski¹, Jonathan Samuel Levy²

¹Dorevitch Pathology, Heidelberg, Vic, Australia; and ²Skin2, St Leonards, NSW, Australia

Syringocystadenocarcinoma papilliferum (SCACP) is an extremely rare malignant adnexal skin tumour with apocrine differentiation. As per our literature search 25 cases of SCACP have been reported so far. It is the malignant counterpart of the more common adnexal skin tumour, syringocystadenoma papilliferum (SCP). They usually arise in a pre-existing long standing stasyringocystadenoma papilliferum, often in the setting of a naevus sebaceous of Jadassohn.¹ A 65-year-old male presented with a history of a scalp lesion that had been growing over a number of years. The H&E stained sections revealed a reticulated lesion composed of markedly atypical appearing cells with basaloid features. Nests of these atypical cells invaded the superficial dermis. The lesion extended to the base of the specimen. Lymphovascular invasion and perineural spread were not observed. CK7 and p63 immunostains revealed focally strong positive staining within the lesion. These findings support the diagnosis of a syringocystadenocarcinoma papilliferum. We also carried out a brief review of literature and found a total of 26 cases of SCACP. These had mean age of 67 years (range 42–93 years), with slight male predominance with male to female ratio of 1.6:1. The most common site was the scalp (half of the cases).

Reference

1. Elder DE, Massi D, Scolyer RA, *et al.*, editors. *WHO Classification of Skin Tumours*. Lyon: International Agency for Research on Cancer, 2018.

COCOON ABDOMEN: BOWELS IN A WEB

Pranav Dorwal¹, Robyn Laurie¹, Kostas Syrrakos²

¹Dorevitch Pathology, Heidelberg, VIC, Australia; and

²Mulgrave Private Hospital, VIC, Australia

Cocoon abdomen is a rare, idiopathic condition characterised by fibrous encapsulation of the bowel segments giving an appearance of a cocoon. The patient may present with variable and non-specific symptoms like abdominal pain, tenderness, vomiting and it sometimes even leads to intestinal obstruction. We hereby report a case of 54-year-old male who presented with symptoms of intestinal obstruction. Apart from the sparing of 60 mm of the resection margin at either ends, most of the segment was encased by the peritoneum giving a 'spider-web' appearance. On opening the intestinal segment, there were multiple areas of kinking of the wall with narrowing of the lumen in the areas of the 'cocooning'. No obvious mass lesion was identified either on the mucosal aspect or in the mesentery. Microscopic examination revealed that multiple loops of small bowel were tethered together by a loose, oedematous fibrovascular membrane. Sclerosing encapsulating peritonitis (SEP) is an extremely rare clinical entity and was first described by Foo *et al.* in 1978,¹ in an adolescent girl, which is reported to be the most common age group. Our case was found in a middle-aged male, which differs in both age group and sex of the common incidence of this entity. The aetiology of this condition remains unknown; however there have been cases where the patients have had a prior abdominal surgery, abdominal tuberculosis, liver transplant or recurrent peritonitis.²

References

1. Foo KT, Ng KC, Rauff A, *et al.* Unusual small intestinal obstruction in adolescent girls: The abdominal cocoon. *Br J Surg* 1978; 65: 427–30.
2. Stanley MM, Reyes CV, Greenlee HB, *et al.* Peritoneal fibrosis in cirrhotics treated with peritoneovenous shunting for ascites. *Digest Dis Sci* 1996; 41: 571–7.

A SPINDLE CELL LESION: CASE REPORT OF A RECTAL GIST PRESENTING A PROSTATIC TUMOUR

Miriam Fewtrell, Mudiwa Muronda

Department of Anatomical Pathology, South Western Area Pathology Service, Liverpool Hospital, Sydney, NSW, Australia

Gastrointestinal stromal tumour (GIST) most commonly occurs in the stomach, followed by the small bowel and uncommonly the colorectum. GISTs can be composed of spindle or epithelioid cells, with most cases characterised by a gain of function mutation of the KIT gene. This case study reports a patient presenting with urinary retention due to a prostatic mass which was biopsied. The histology was of a mitotically active spindle cell lesion with necrosis. Positive staining for CD117 and DOG1 confirmed the diagnosis of GIST. Subsequent imaging demonstrated a rectal origin of the tumour.