reactive polymorphous inflammatory background – features characteristic of AITL. Careful assessment also identified a second neoplastic population of large centroblast-like B lymphocytes. The neoplastic T cells were immunopositive for pan-T-cell antigens CD43, CD2, CD3 and CD5; and for markers of T follicular helper cell immunophenotype PD1 and CXCL13. The large centroblast-like B cells showed immunoreactivity to pan-B-cell antigens CD20, CD79a, PAX5 and OCT2, and to EBER signifying EBV-infection. CLs are unusual diagnoses but should be considered when there are clinical, morphological or immunophenotypical inconsistencies. In addition, the particular combination of lymphomas present impacts treatment decisions.

AN UNUSUAL BIPHASIC PULMONARY LESION: LATE METASTASIS FROM A MANDIBULAR AMELOBLASTOMA

Thomas de Hoog, Peter Jessup

Department of Anatomical Pathology, Royal Hobart Hospital, Hobart, Tas, Australia

Ameloblastoma is mostly a benign odontogenic tumour most frequently arising within the mandible and maxilla. Although ameloblastoma can be locally aggressive and prone to local recurrence, metastasis is a rare (and usually late) complication. We report a case of a metastatic ameloblastoma to the lung following an original presentation with a traumatic jaw fracture through what was thought to be a mandibular odontogenic cyst in 1988. The definitive diagnosis of ameloblastoma was not reached until a local recurrence occurred ten years later; this was managed with hemimandibulectomy and post-operative radiotherapy. A single lung lesion was identified incidentally 33 years after the original presentation: it demonstrated typical morphological features of ameloblastoma and FOXP1 immunostaining was confirmatory. Recent advances in understanding the genetic pathogenesis of ameloblastoma demonstrate frequent *BRAF*^{V600E} mutations as well as mutations in other MAP kinase pathway genes such as *KRAS*, *NRAS*, *HRAS* and *FGFR2*.¹ Several case reports have indicated promising results with *BRAF* inhibitor therapy in metastatic ameloblastoma.²

References

1. Brown N, Betz B. Ameloblastoma: a review of recent molecular pathogenetic discoveries. *Biomark Cancer* 2015; 7 (Suppl 2): 19–24.
2. González-González R, López-Verdín S, Lavalle-Carrasco J, *et al.* Current concepts in ameloblastoma-targeted therapies in B-raf proto-oncogene serine/threonine kinase V600E mutation: Systematic review. *World J Clin Oncol* 2020; 11: 31–42.

NEPHROPATHY IN FABRY DISEASE

Marsa Hosseinzadeh¹, Gopala Rangan², Michel Tchan³, Carol Robinson¹, Seethalakshmi Viswanathan¹

¹Department of Anatomical Pathology, ICPMR, Westmead Hospital, NSW, Australia; ²Department of Renal Medicine, Westmead Hospital, NSW, Australia; and ³Department of Genetic Medicine, Westmead Hospital, NSW, Australia

Fabry disease (FD) is a rare, progressive, X-linked multisystem lysosomal storage disorder caused by the partial or total deficiency of the lysosomal enzyme α -galactosidase A.

Herein, we report a case of a 24-year-old man with genetically proven Fabry disease who presented with angiokeratoma, lymphadenoma, feet paraesthesia and non-specific gastrointestinal symptoms including abdominal bloating and irregular bowel motions.

He had a family history of FD in his mother (carrier) and older brother who was receiving Enzyme Replacement Therapy (ERT). The renal biopsy was organised to assess the extent of renal involvement and eligibility for (ERT).

On light microscopy, the glomeruli showed characteristic podocytes with foamy vacuolated cytoplasm. There were mild background chronic changes and mild arteriosclerosis.

The electron microscopic examination revealed characteristic myeloid bodies in podocytes, interstitium, tubular epithelial cells, endothelial cells of peritubular capillaries and the arterioles. These myeloid bodies showed concentrically arranged lamellae of electron dense material.

The diagnosis of FD is often delayed due to its non-specific symptoms in the initial stage. It is recommended that ERT should be initiated as early as possible, and this underlies the importance of early detection of Fabry disease.

References

1. Wani MM, Khan I, Bhat RA, *et al.* Fabry's disease: case series and review of literature. *Ann Med Health Sci Res* 2016; 6: 193–7.
2. Duro G, Zizzo C, Cammarata G, *et al.* Mutations in the GLA gene and LysoGb3: is it really Anderson-Fabry disease? *Int J Mol Sci* 2018; 19: 3726.
3. Alegra T, Vairo F, de Souza MV, *et al.* Enzyme replacement therapy for Fabry disease: A systematic review and meta-analysis. *Genet Mol Biol* 2012; 35 (4 Suppl): 947–54.

PRIMARY OVARIAN DIFFUSE LARGE B-CELL LYMPHOMA

Daniel Hussey, Sophie Bittinger

Anatomical Pathology Department, The Royal Women's Hospital, Parkville, Vic, Australia

The ovary is a common site for lymphoma in disseminated disease, but primary ovarian lymphoma is exceedingly rare.¹ Primary non-Hodgkin's lymphomas constitute 1.5% of ovarian malignancies.¹ The diagnosis carries a poor prognosis with a 5-year survival of 0 to 36%.² This case study reports a 27-year-old female with presenting with abdominal pain and urinary symptoms. Investigations found a large right adnexal mass which was removed laparoscopically. Microscopy showed replacement of the ovary by discohesive cells with high N:C ratio and scant cytoplasm. The cells had enlarged angular nuclei, clumped chromatin and small nucleoli. Immunohistochemistry for CD45, CD20, bcl-2, bcl-6, MUM-1 and PAX5 was positive. Flow cytometry showed a large clonal CD5- CD10- B cell population. These findings were consistent with a diffuse large B-cell lymphoma. Although rare the diagnosis primary ovarian lymphoma should be considered as a differential in round blue cell malignancies of the ovary. Significant mortality means early diagnosis is key for a favourable outcome.

References

1. Vang R, Medeiros L, Warnke R, *et al.* Ovarian non-Hodgkin's lymphoma: a clinicopathologic study of eight primary cases. *Mod Pathol* 2001; 14: 1093–9.
2. Hu R, Miao M, Zhang R, *et al.* Ovary involvement of diffuse large B-cell lymphoma. *Am J Case Rep* 2012; 13: 96–8.