Fig. 2 Pseudoacini positive on immunostaining for (A) CD45, (B) CD20, (C) CD10, (D) bcl2 and (E) bcl6. (F) Ki-67 staining is higher in the clusters of large cells than in the background smaller lymphocytes (bone marrow trephine).

index was low, but staining generally corresponded to the clusters of large cells (Fig. 2F). The background small lymphocytes had an immunophenotype consistent with FL, these cells being CD20+, CD5–, CD10+ and bcl2+. An excisional lymph node biopsy was subsequently performed, confirming a diagnosis of DLBCL, germinal centre type, on a background of FL. The patient proceeded to receive chemo-immunotherapy and obtained a complete metabolic remission.

Lymphomas can display unusual cohesive growth patterns. In the case of anaplastic large cell lymphoma,³ the striking intrasinusoidal infiltration and perivascular homing of the lymphoma cells may be explained by the expression of clusterin.⁴ Likewise, the formation of rosettes in a manner akin to neuroepithelial and neuroendocrine tumours^{5–7} has been described in both B- and T-cell lymphomas at nodal and extra-nodal sites.^{8–11} The acellular fibrillary material that produces such rosettes' central lumen is believed to result from the long interdigitating cytoplasmic processes of lymphoma cells with microvilli or from adjacent histiocytes.⁷

Herein, we report a unique manifestation of tumour architecture leading to the formation of discrete large cell aggregates on a background of FL, separate from vascular sinuses and without rosette formation. This peculiar motif is rare; while it has been described in a case of DLBCL affecting a lymph node,¹² it has not been reported in the bone marrow. In our case, this feature was observed only in the bone marrow and not in the lymph node, even though large cell transformation was present at both sites. The histogenesis of our case is uncertain. Conceivably, the architecture depicted here may be secondary to surface glycoproteins mediating cell-to-cell adhesion amongst the larger cells. Alternatively, the nests of larger cells may represent the initial seeds of large cell transformation which, given time, may have proliferated to infiltrate the marrow diffusely.

Lymphoma histology can be protean, and our case illustrates a unique histological manifestation of DLBCL.

Without the benefit of the patient's past clinical history and immunohistochemistry, this case may have been misdiagnosed as metastatic carcinoma. Our case is intended to serve as a cautious lesson on the mutable nature of lymphoma histology and the potential for diagnostic confusion.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest or sources of funding to disclose.

Matthew J. Rees¹, Chris Hogan¹, Tineke Fancourt², Wai Khoon Ho¹

¹Department of Laboratory Haematology, Austin Health, Melbourne, Vic, Australia; ²Department of Anatomical Pathology, Austin Health, Melbourne, Vic, Australia

Contact Dr Matthew James Rees.

E-mail: Matthew.Rees@austin.org.au

1. Wagner-Johnston ND, Link BK, Byrtek M, *et al.* Outcomes of transformed follicular lymphoma in the modern era: a report from the National LymphoCare Study (NLCS). *Blood* 2015; 126: 851–7.
2. Smith S. Transformed lymphoma: what should I do now? *Hematology* 2020; 2020: 306–11.
3. Montes-Mojarro IA, Steinhilber J, Bonzheim I, *et al.* The pathological spectrum of systemic anaplastic large cell lymphoma (ALCL). *Cancers (Basel)* 2018; 10: 107.
4. Jaffe ES. Anaplastic large cell lymphoma: the shifting sands of diagnostic hematopathology. *Mod Pathol* 2001; 14: 219–28.
5. Frizzera G, Gajl-Peczalska K, Sibley RK, *et al.* Rosette formation in malignant lymphoma. *Am J Pathol* 1985; 119: 351–6.
6. Alakeel F, DePalma L. Follicular lymphoma with rosette formation with an exclusive diffuse pattern. *Virchows Arch* 2017; 470: 231–3.
7. Koo CH, Shin SS, Bracho F, *et al.* Rosette-forming non-Hodgkin's lymphomas. *Histopathology* 1996; 29: 557–63.
8. Rekhi B, Gulia A. Rosette forming" primary high-grade B-cell lymphoma, presenting as a scapular mass, masquerading Ewing sarcoma: a rare case with diagnostic implications. *Indian J Pathol Microbiol* 2021; 64: 617–9.
9. Sioutos N, Kadin ME. Perivascular ki-1 + lesions. *Int J Hematol* 1992; 55: 275–9.
10. Goteri G, Costagliola A, Tasseti A, *et al.* Diffuse large B-cell lymphoma with Homer-Wright rosettes, sinusoidal growth pattern, and CD30 expression: a possible overlap between microvillous lymphomas and sinusoidal CD30-positive large B-cell lymphomas. *Pathol Res Pract* 2009; 205: 279–82.
11. Lai R, Medeiros LJ, Dabbagh L, *et al.* Sinusoidal CD30-positive large B-cell lymphoma: a morphologic mimic of anaplastic large cell lymphoma. *Mod Pathol* 2000; 13: 223–8.
12. Mok Y, Goh GH, Wang S. Unusual cohesive growth pattern of diffuse large B-cell lymphoma mimicking metastatic carcinoma. *Br J Haematol* 2017; 176: 850.

DOI: <https://doi.org/10.1016/j.pathol.2021.10.013>

Validation of *EGFR* mutation testing on cytological smears of lung cancer using the Idylla platform

To the Editor,

Patients with stage IV lung cancer have a poor prognosis with only 19% of patients surviving 12 months in Australia.¹ However, for patients with tumours containing specific molecular alterations (such as activating mutations in *EGFR*,

ROS1 and *ALK* genes) targeted tyrosine kinase inhibitor (TKI) therapy can provide significantly longer survival. In clinical studies, approximately 10–20% of biopsies are inadequate for *EGFR* mutation testing.²

Endobronchial ultrasound transbronchial needle aspiration (EBUS-TBNA) sampling of lymph nodes is an important technique in diagnosing and staging of patients with lung cancer. In this study, we sought to validate the use of Diff-Quik stained slides from EBUS-TBNA procedures as input material for *EGFR* mutation testing using the Idylla (Bio-cartis, Belgium) platform.

Archived Diff-Quik stained cytology smears (between 2014–2020) containing tumour cells from EBUS-TBNA were obtained from 54 patients with lung adenocarcinoma with known *EGFR* mutation status. In 33 patients, *EGFR* mutation status was determined from a formalin fixed, paraffin embedded (FFPE) cell-block specimen at the time of EBUS-TBNA. In the other 21 patients, *EGFR* mutation status was determined from a different FFPE specimen, either prior or after the EBUS-TBNA procedure. In 7/54, the *EGFR* mutation status had been determined using the Idylla *EGFR* assay, 15 were by next generation sequencing (NGS), 30 by pyrosequencing, one by competitive allele-specific TaqMan polymerase chain reaction (PCR) and one by Therascreen RGQ PCR. The use of patient material and clinical information was approved by the Mater Misericordiae Ltd Human Research Ethics Committee [review reference: HREC/MML/62296(V3)].

Diff-Quik stained slides were scanned at 40× using a CS120 slide scanner (Olympus, Japan) to preserve the medical record and allow for tumour cellularity assessment. Two pathologists estimated the total smear cellularity using thresholds as previously described.³ The tumour fraction was estimated using quartiles: <25%, 25–50%, 50–75% and 75–100%. Stained slides were deconverslipped by immersing in xylene for up to 3 days. Slides were then immersed in 50% xylene/alcohol for 5 min and absolute alcohol for 5 min. The slide from Patient 1 was not destained; this case without known *EGFR* mutations was used to determine whether autofluorescence was detectable. Slides from Patients 2–54 were destained by dipping in 0.2% acid alcohol until the Diff-Quik stain was visibly removed (approximately 20 dips). Slides were then immersed in distilled water for 30 s and air dried. Cells were scraped from the slide using a scalpel onto dampened sterile circular filter paper and placed into the cartridge. No enrichment or targeted sampling was attempted.

The internal *EGFR* quantification cycle (Cq) was compared between cases using unpaired two-tailed t-tests. In patient samples that had matched Idylla *EGFR* mutation assays performed on their diagnostic sample and Diff-Quik stained slides, the Wilcoxon matched pairs signed rank test was used to determine significant differences in internal *EGFR* Cq values. Significance was determined if $p \leq 0.05$.

A previous study has identified background fluorescence in the Idylla system's channel 2.⁴ We performed one assay (Patient 1) without first destaining and saw high background fluorescence compared to assays run with destaining (Fig. 1, Patient 1). The instrument recorded approximately 600 relative fluorescence units (RFU) in channel 2 without destaining, which was reduced to approximately 20 RFU with destaining (Fig. 1, Patient 2). Graphs from Patient 4 with an L858R mutation show that fluorescence increased to just over

Patient 1 - no *EGFR* mutations, no destaining

Patient 2 - no *EGFR* mutations, with destaining

Patient 4 - L858R mutation, with destaining

Fig. 1 Destaining Diff-Quik stained slides reduces background fluorescence in channel 2, responsible for detection of L858R mutations in *EGFR*. Amplification curves for channel 2 for Patients 1, 2 and 4. The sample from Patient 1 without detectable *EGFR* mutations shows high background fluorescence without destaining. The sample from Patient 2 without detectable *EGFR* mutations shows much lower background fluorescence in channel 2 with destaining. Patient 4 has an L858R mutation in *EGFR* represented by a positive amplification curve (ChD).

40 RFU in channel 2 when the mutation was present, which was typical for detection of this mutation.

Overall, 54 patients were included in the validation study. Of these, 17 were known to have a mutation in *EGFR* (31.4%). All patients with a known mutation in *EGFR* were also detected from the Diff-Quik stained smear (Table 1). There were no false positives.

In two cases (Patients 6 and 19), the original diagnostic result indicated co-occurring mutations in *EGFR* (exon 19 deletion and T790M), of which only the exon 19 deletion was detected in the Diff-Quik stained smear slide.

In Patient 19, the diagnostic biopsy was obtained 12 months after the EBUS-TBNA Diff-Quik stained sample and

Table 1 Results for patients included in the validation study ($n=54$)

Patient no.	Diagnostic sample	Diagnostic method	Diagnostic EGFR result	DQ cellularity	DQ tumour proportion	DQ EGFR result
1	Biopsy	NGS	NMD	Moderate	25–50%	NMD
2	Cell block	Idylla	NMD	Moderate	25–50%	NMD
3	Biopsy	Idylla	Ex19del	Moderate	>75%	Ex19del
4	Biopsy	Idylla	L858R	Low	<25%	L858R
5	Cell block	Idylla	L858R	Low	<25%	L858R
6	Biopsy	Pyrosequencing	Ex19del, T790M	Moderate	<25%	Ex19del ^a
7	Cell block	NGS	Ex20ins	High	25–50%	Ex20ins
8	Biopsy	Idylla	NMD	Moderate	50–75%	NMD
9	Cell block	NGS	NMD	Moderate	50–75%	NMD
10	Cell block	NGS	NMD	Moderate	25–50%	NMD
11	Cell block	Idylla	NMD	Moderate	25–50%	NMD
12	Biopsy	Pyrosequencing	Ex19del	High	<25%	Ex19del
13	Cell block	Idylla	Ex19del	Moderate	<25%	Ex19del
14	Cell block	NGS	Ex20ins	Moderate	50–75%	Ex20ins
15	Cell block	NGS	L858R	Moderate	>75%	L858R
16	Cell block	Pyrosequencing	L858R	High	>75%	L858R
17	Biopsy	Pyrosequencing	Ex19del	Moderate	25–50%	Ex19del
18	Cell block	Pyrosequencing	Ex19del, T790M	Moderate	25–50%	Ex19del, T790M
19	Biopsy	Pyrosequencing	Ex19del, T790M	Moderate	25–50%	Ex19del ^a
20	Cell block	NGS	Ex19del	Moderate	25–50%	Ex19del
21	Biopsy	NGS	Ex19del, T790M	Moderate	25–50%	Ex19del, T790M
22	Cell block	Pyrosequencing	Ex19del	Moderate	50–75%	Ex19del
23	Cell block	Pyrosequencing	NMD	High	<25%	NMD
24	Biopsy	Pyrosequencing	NMD	Moderate	>75%	NMD
25	Cell block	Pyrosequencing	NMD	Moderate	25–50%	NMD
26	Cell block	Pyrosequencing	NMD	Moderate	25–50%	NMD
27	Biopsy	TaqMan	L861Q	Moderate	50–75%	L861Q
28	Biopsy	Pyrosequencing	NMD	High	>75%	NMD
29	Cell block	Pyrosequencing	NMD	Moderate	25–50%	NMD
30	Biopsy	Pyrosequencing	NMD	Moderate	50–75%	NMD
31	Cell block	Pyrosequencing	NMD	High	>75%	NMD
32	Biopsy	Pyrosequencing	NMD	Low	>75%	NMD
33	Cell block	Pyrosequencing	NMD	High	>75%	NMD
34	Cell block	Pyrosequencing	NMD	Moderate	>75%	NMD
35	Cell block	Pyrosequencing	NMD	Moderate	25–50%	NMD
36	Biopsy	Therascreen	NMD	High	25–50%	NMD
37	Biopsy	Pyrosequencing	NMD	High	50–75%	NMD
38	Cell block	NGS	NMD	Moderate	25–50%	NMD
39	Cell block	Pyrosequencing	NMD	Moderate	50–75%	NMD
40	Cell block	Pyrosequencing	NMD	Moderate	>75%	NMD
41	Biopsy	NGS	NMD	Moderate	>75%	NMD
42	Cell block	Pyrosequencing	NMD	High	25–50%	NMD
43	Cell block	Pyrosequencing	NMD	High	>75%	NMD
44	Cell block	NGS	NMD	Moderate	50–75%	NMD
45	Cell block	Pyrosequencing	NMD	Moderate	25–50%	NMD
46	Cell block	Pyrosequencing	NMD	High	>75%	NMD
47	Biopsy	Pyrosequencing	NMD	Moderate	>75%	NMD
48	Cell block	NGS	NMD	Moderate	>75%	NMD
49	Cell block	Pyrosequencing	NMD	High	>75%	NMD
50	Cell block	NGS	NMD	Moderate	50–75%	Error ^b
51	Cell block	NGS	NMD	Moderate	25–50%	NMD
52	Cell block	Pyrosequencing	NMD	Moderate	50–75%	NMD
53	Cell block	Pyrosequencing	NMD	High	50–75%	NMD
54	Cell block	NGS	NMD	Moderate	>75%	NMD

DQ, Diff-Quik; NGS, next generation sequencing; NMD, no mutation detected.

54 patients with lung adenocarcinoma with known *EGFR* mutation status had DQ stained slides from EBUS-TBNA used as input material for the Idylla *EGFR* assay. *EGFR* mutation status was determined from diagnostic samples, either small biopsy samples or EBUS-TBNA samples made into a formalin-fixed, paraffin-embedded cell block. The method used diagnostically was by NGS, Idylla *EGFR* assay, pyrosequencing (Qiagen), Competitive Allele Specific Taqman PCR or Therascreen RGQ PCR (Qiagen). Total cellularity was estimated to be low (<1000 cells), moderate (1000–5000 cells) or high (>5000 cells). Tumour proportion estimates are expressed as a percentage of total cellularity.

^a Potentially discrepant results (see text);

^b Test failure.

the patient had previously been treated with gefitinib. The cytology sample had been obtained prior to the development of resistance and therefore it is not expected that this sample would contain the T790M mutation.

In Patient 6, the EBUS-TBNA sample had been obtained approximately 24 months after the diagnostic biopsy. The diagnostic small biopsy had been taken from a pleural based nodule, whilst the cytology sample had been taken from a station 7 lymph node. We identified an additional Diff-Quik stained slide obtained from the same procedure and extracted DNA for NGS. The only mutation in *EGFR* with a variant allele fraction of $>10\%$ was an exon 19 deletion

(c2235_2249del) with no evidence of T790M. This is likely a case of heterogeneity of T790M positive clones.⁵

One sample (patient 50) was associated with cartridge failure and was unable to provide a result. An accurate *EGFR* mutation was detected even in samples with low cellularity or low tumour fraction (Fig. 2A).

Increasing smear cellularity was associated with lower Cq for the internal *EGFR* control, consistent with increased input of amplifiable DNA (Fig. 2B). Diff-Quik stained slides with high cellularity had significantly lower Cq values for the internal *EGFR* control than moderate or low cellularity smears. In paired tumour samples, where the Idylla *EGFR*

Fig. 2 Representative Diff-Quik smears from patients with detectable *EGFR* mutations and a comparison of Cq values for internal *EGFR* control. (A) Upper panels: Diff-Quik stained slide from Patient 4 with low cellularity and $<25\%$ tumour fraction. Area of high power view indicated by box. Lower panels: Patient 6 with moderate overall cellularity but low (well less than 25%) tumour fraction. (B) Cq values for low, moderate and high cellularity smears; lines represent mean and standard deviation. (C) For six patients with matched formalin-fixed paraffin-embedded diagnostic samples, the reference Cq value was compared to the Diff-Quik stained slide. ns, not significant; * $p \leq 0.05$; ** $p \leq 0.005$.

mutation test was performed on the formalin-fixed paraffin-embedded samples, the internal *EGFR* control had a significantly lower Cq in the Diff-Quik stained smears than formalin-fixed samples (Fig. 2C).

A recent study investigated the use of the Idylla system in 301 samples of lung cancer, 13 of which were scraped from Diff-Quik stained slides without destaining.⁶ Two of these 13 cases were not detected by the Idylla software, one of which had an L858R mutation, the other a deletion in Exon19 (del15), although the authors noted that they had mutation curves on manual review of the unfiltered data. It is likely that in the presence of high background fluorescence Diff-Quik stained slides could lead to false negatives, particularly in channel 2 responsible for the detection of L858R. It is noteworthy that we were able to detect L858R in all four of our cases using our destaining protocol, even though these cases had both low cellularity and low tumour proportion.

In cell line dilution studies, a total *EGFR* Cq at less than or equal to 23 allows for the detection of exon 19 deletions, L858R and T790M mutations when their variant allele frequency is 5% (10% cellularity).⁶ Eleven of our 53 successfully run samples had *EGFR* Cq values above 23, although all paired samples tested exhibited lower Cq values than their matched FFPE samples. Our findings suggest that Diff-Quik stained cytology smears is a preferable input material to small biopsies.

Two cases in our series failed to detect concurrent T790M mutations, one (Patient 19) due to the patient not developing the resistance mutation at the time the EBUS-TBNA sample was taken, the other (Patient 6) potentially due to heterogeneity of metastatic disease sites. Reduced sensitivity for the T790M mutations have been described on the Idylla platform, associated with higher *EGFR* control Cq values and Q787Q polymorphism in cis.⁷ This reduced sensitivity for T790M as well as the absence of primers for other mutations including C797S and G724S make the Idylla platform unsuitable for testing in the setting of TKI resistance.⁶ It is notable that we were able to detect concurrent T790M mutations in two other samples (Patients 18 and 21). Excessive input DNA has been reported to be associated with false positives.⁸ Although we did not detect any false positives in this study, a manual review of amplification curves is performed on each case routinely in our laboratory.⁶

During the EBUS-TBNA procedure it is not uncommon for the pathologist to report tumour cells during rapid on-site evaluation (ROSE) but subsequent needle passes fail to yield sufficient diagnostic material for a cell block. Insufficient material for mutation testing occurs in 5–25% of patients.³ The routine use of NGS panels⁹ can lead to a delay in providing results and NGS approaches can fail in up to 10% of cases sampled by EBUS-TBNA.¹⁰ By using the Diff-Quik stained slide, laboratories could rapidly identify the 15–40% of patients with adenocarcinoma with *EGFR* mutations without using cell block or biopsy material which may be used for subsequent comprehensive NGS profiling. There are significant cost benefits to this approach, including a reduced number of additional procedures due to inadequate sampling and assay failure.

In conclusion, this study demonstrates that utilising the Idylla *EGFR* mutation assay on destained Diff-Quik stained slides obtained from EBUS-TBNA procedures is 100% concordant with current laboratory based methods for

mutation testing, even in low cellularity smears without enrichment. The control *EGFR* Cq values suggests that Diff-Quik stained slides are superior to FFPE material for this assay. By utilising this method in the management of patients with lung cancer, clinicians can obtain a rapid same day result for *EGFR* mutation status, reserving FFPE cell block and biopsy material for immunohistochemistry, fluorescence *in situ* hybridisation and NGS and reducing the need for further diagnostic procedures with associated risks.

Acknowledgements: We thank Felicity Murphy and Bree Gall who provided clinical advice and scientific input.

Conflicts of interest and sources of funding: The Idylla *EGFR* mutation test cartridges were supplied by Biocartis. Biocartis had no role in study design, analysis and interpretation of the data, in the writing of the report or in the decision to submit the paper for publication. The authors declare no other conflicts of interest.

Cameron Snell^{1,2}, Sam McManus¹, Kelly Robinson¹, Wendy Lewis¹, Samantha Arandelovic¹, Rohan Lourie¹, James Harraway¹

¹Mater Pathology, Mater Hospital Brisbane, South Brisbane, Qld, Australia; ²Cancer Pathology Research Group, Mater Research Institute, University of Queensland, Translational Research Institute, Brisbane, Qld, Australia

Contact Dr Cameron Snell.

E-mail: Cameron.snell@petermac.org

1. Cancer Australia, National Cancer Control Indicators. Relative Survival by Stage at Diagnosis (Lung Cancer). 1 Apr 2019; cited Apr 2021. <https://nccci.cancer australia.gov.au/outcomes/relative-survival-rate/relative-survival-stage-diagnosis-lung-cancer>
2. Douillard JY, Ostoros G, Cobo M, *et al.* Gefitinib treatment in *EGFR* mutated caucasian NSCLC: circulating-free tumor DNA as a surrogate for determination of *EGFR* status. *J Thorac Oncol* 2014; 9: 1345–53.
3. Fielding D, Dalley AJ, Bashirzadeh F, *et al.* Diff-Quik cytology smears from endobronchial ultrasound transbronchial needle aspiration lymph node specimens as a source of DNA for next-generation sequencing instead of cell blocks. *Respiration* 2019; 97: 525–39.
4. De Luca C, Conticelli F, Leone A, *et al.* Is the Idylla *EGFR* Mutation Assay feasible on archival stained cytological smears? A pilot study. *J Clin Pathol* 2019; 72: 609–14.
5. Ichihara E, Lovly CM. Shades of T790M: intratumor heterogeneity in *EGFR*-mutant lung cancer. *Cancer Discov* 2015; 5: 694–6.
6. Arcila ME, Yang SR, Momeni A, *et al.* Ultrarapid *EGFR* mutation screening followed by comprehensive next-generation sequencing: a feasible, informative approach for lung carcinoma cytology specimens with a high success rate. *JTO Clin Res Rep* 2020; 1: 100077.
7. Lee E, Jones V, Topkas E, *et al.* Reduced sensitivity for *EGFR* T790M mutations using the Idylla *EGFR* mutation test. *J Clin Pathol* 2021; 74: 43–7.
8. Caputo A, D'Ardia A, Sabbatino F, *et al.* Testing *EGFR* with Idylla on cytological specimens of lung cancer: a review. *Int J Mol Sci* 2021; 22: 4852.
9. Lindeman NI, Cagle PT, Aisner DL, *et al.* Updated molecular testing guideline for the selection of lung cancer patients for treatment with targeted tyrosine kinase inhibitors: guideline from the College of American Pathologists, the International Association for the Study of Lung Cancer, and the Association for Molecular Pathology. *Arch Pathol Lab Med* 2018; 142: 321–46.
10. Turner SR, Buonocore D, Desmeules P, *et al.* Feasibility of endobronchial ultrasound transbronchial needle aspiration for massively parallel next-generation sequencing in thoracic cancer patients. *Lung Cancer* 2018; 119: 85–90.

DOI: <https://doi.org/10.1016/j.pathol.2021.10.006>