from the learning that is a consequence of receiving MALDI-TOF MS results back to either confirm or refute the scientists' suspected identification based on the colony morphology. In addition, staff described a loss of professional satisfaction when feeling 'disconnected' from the process of following a specimen through from beginning to the end.

Importantly, with respect to quality improvement, staff described that the process of following up one's own work was more likely to reduce errors since receiving back an unexpected MALDI-TOF MS result was a trigger to consider whether the wrong colony might have been spotted onto the MALDI-TOF MS target, whether the culture was mixed or whether the inoculum was misplaced onto an incorrect spot on the MALDI-TOF MS target as has been described in Public Health England UK Microbiology standards.¹² This impression was supported by findings from this survey conducted after the change of TLA workflow. Further studies could support our findings by including error audits before and after TLA workflow changes.

Limitations of our study include the assessment of contributing factors to job satisfaction particularly in question two, as it is not clear if the reason for the change in job satisfaction was due to the Kiestra TLA or due to other factors or changes in the laboratory that occurred during the same time period. Given that other responses (to questions three, four and five) indicate Kiestra TLA workflow changes were mostly associated with 'much better' outcomes, this implies that lack of change in job satisfaction overall is likely due to factors unrelated to the TLA. In addition, the wording of question one may have selected more experienced staff to respond to the survey: since the impact of laboratory changes may be different for more recently qualified staff as compared to experienced staff, further studies are required to explore this possibility. Further studies are warranted to examine the objective impact of TLA workflow changes on quality metric such as turnaround time and error rates.

In conclusion, after implementing a change to Kiestra TLA workflow we found that job satisfaction improved for the vast majority of the staff. This change was associated with a perceived positive impact on continuing professional development and education and on quality and reduction of errors compared to the unidirectional TLA workflow. Microbiology laboratory managers should be aware of the impact of various TLA workflows on their staff as this is likely to contribute to quality improvement.

Acknowledgements: We thank George Streitberg from Monash Health Biochemistry for reviewing and providing advice on the design and wording of the survey questions.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest to disclose.

Maryza Graham^{1,2,3}, Emma Lee¹, Richard Streitberg¹, John Hamblin¹, Beena Kumar⁴, Tony M. Korman^{1,2,3}

¹Department of Microbiology, Monash Health Pathology, Monash Health, Clayton, Vic, Australia; ²Monash Infectious Diseases, Monash Health, Clayton, Vic, Australia; ³Faculty of Medicine, Nursing and Health Sciences, Monash University, Clayton, Vic, Australia; ⁴Monash Health Pathology, Monash Health, Clayton, Vic, Australia

Contact Dr Maryza Graham.

E-mail: Maryza.Graham@monashhealth.org

1. Froment P, Marchandin H, Vande Perre P, *et al.* Automated versus manual sample inoculations in routine clinical microbiology: a performance evaluation of the fully automated Inoqula instrument. *J Clin Microbiol* 2014; 52: 796–802.
2. Mutters NT, Hodiamont CJ, de Jong MD, *et al.* Performance of Kiestra total laboratory automation combined with MS in clinical microbiology practice. *Ann Lab Med* 2014; 34: 111–7.
3. Croxatto A, Dijkstra K, Prod'homme G, *et al.* Comparison of inoculation with the Inoqula and WASP automated systems with manual inoculation. *J Clin Microbiol* 2015; 53: 2298–307.
4. Strauss S, Bourbeau PP. Impact of introduction of the BD Kiestra Inoqula on urine culture results in a hospital clinical microbiology laboratory. *J Clin Microbiol* 2015; 53: 1736–40.
5. Jacot D, Sarton-Lohéac G, Coste AT, *et al.* Performance evaluation of the becton dickinson Kiestra™ IdentifA/SusceptA. *Clin Microbiol Infect* 2021 Aug; 27: 1167.e9–17.
6. Croxatto A, Prod'homme G, Faverjon F, *et al.* Laboratory automation in clinical bacteriology: what system to choose? *Clin Microbiol Infect* 2016; 22: 217–35.
7. Graham M, Tilson L, Streitberg R, *et al.* Improved standardization and potential for shortened time to results with BD Kiestra™ total laboratory automation of early urine cultures: a prospective comparison with manual processing. *Diagn Microbiol Infect Dis* 2016; 86: 1–4.
8. Yarbrough ML, Lainhart W, McMullen AR, *et al.* Impact of total laboratory automation on workflow and specimen processing time for culture of urine specimens. *Eur J Clin Microbiol Infect Dis* 2018; 37: 2405–11.
9. Culbreath K, Piwonka H, Korver J, *et al.* Benefits derived from full laboratory automation in microbiology: a tale of four laboratories. *J Clin Microbiol* 2021; 59: e01969–20.
10. Dauwalder O, Landrieu L, Laurent F, *et al.* Does bacteriology laboratory automation reduce time to results and increase quality management? *Clin Microbiol Infect* 2016; 22: 236–43.
11. Klein S, Nurjadi D, Horner S, *et al.* Significant increase in cultivation of *Gardnerella vaginalis*, *Alloscardovia omnicolens*, *Actinotignum schaalii*, and *Actinomyces* spp. in urine samples with total laboratory automation. *Eur J Clin Microbiol Infect Dis* 2018; 37: 1305–11.
12. Public Health England. *UK Standards for Microbiology Investigations: Matrix-Assisted Laser Desorption/Ionisation - Time of Flight Mass Spectrometry (MALDI-TOF MS) Test Procedure*. TP 40 Issue 1.1. London: Public Health England, 2019. <https://www.gov.uk/government/publications/smi-tp-40-maldi-tof-ms-test-procedure>

DOI: <https://doi.org/10.1016/j.pathol.2021.10.010>

Invasive extended-spectrum beta-lactamase (ESBL) related infection rates are not influenced by international travel restrictions

To the Editor,

Escalating antimicrobial resistance (AMR) is recognised by the World Health Organization and World Economic Forum as a major global health and security threat, with supply of new drugs vastly outstripped by the rate of resistance development. Global rates of extended spectrum beta-lactamase (ESBL) carriage in healthy populations have risen 8-fold to 21% over the last 15 years,¹ and recent estimates suggest that 700,000 deaths per year are attributable to AMR worldwide.² Increasing resistance profiles amongst Gram negative bacteria are of particular clinical concern due to the limited repertoire of antibiotics that are effective against these organisms and their ability to readily exchange drug resistance elements through horizontal gene transfer.³ Key factors driving high resistance rates include

widespread travel, variable antibiotic stewardship frameworks, limited community awareness of appropriate antibiotic use, unregulated agricultural application of antimicrobials and health system factors, which all contribute to excessive and poorly targeted antibiotic use.¹

Rates of drug resistance are highest in low income settings, including regions of South and South East Asia.¹ AMR carriage is rising at an alarming rate in high income countries such as Australia also.⁴ Studies have suggested that international travel to high ESBL prevalence countries may be a risk factor for ESBL carriage in high income countries.⁵ Indeed European reports have highlighted high rates of ESBL carriage in stool after return from travel to Asia, with prevalence decreasing almost to baseline over the subsequent 12 months.⁶

Public health measures instituted in the context of the COVID-19 pandemic have included extreme limitation on international travel, introduced in Australia on 20 March 2020. Since that time, incoming passenger numbers reduced by two orders of magnitude from an average of 1.8 million people/month (prior to March 2020) to 24,000 people/month (between April 2020 and March 2021).⁷ This prolonged period with minimal influx of travellers provides a unique opportunity to analyse the impact of international travel on the prevalence of community acquired ESBL-related disease.

Invasive ESBL-related infections carry high morbidity and mortality for the individual and substantial health care burden due to limited treatment repertoires and reliance on antibiotics that often carry high toxicities. This study focuses on invasive community acquired ESBL disease as a critical medical issue in the hospital setting, while recognising this is the tip of the iceberg of overall AMR carriage. Known circulation of ESBLs in long term care facilities and possible hospital acquired infections are expected to follow different epidemiological patterns,⁸ and were excluded from this report.

The proportion of community acquired sterile site Enterobacteriaceae isolates expressing an ESBL phenotype at an urban tertiary hospital in Sydney was analysed over the 12 month period following international travel restrictions (April 2020–March 2021) in comparison to the prior 5 year period (April 2015–March 2020). Infections in older persons (>70 years) and long term care facility residents and those acquired more than 48 hours after hospital admission were excluded. Similarly, recurrent infections due to an organism with the same genotype were excluded as these were likely to be representative of persistent gastrointestinal carriage with translocation or re-inoculation to sterile sites rather than a new exogenous community acquired infection. ESBL production was determined phenotypically utilising double-disc diffusion confirmation testing using cefotaxime (30 µg), ceftazidime (30 µg) and cefepime (30 µg) as indicator cephalosporins, with and without supplemental clavulanic acid (10 µg) according to standard Clinical and Laboratory Standards Institute (CLSI) methodologies.⁹ Targeted nucleic acid amplification testing (NAAT) for CTX-M, IMP-4 and pan-IMP genotypes using an in-house assay was conducted on all isolates using established protocols¹⁰ to determine the dominant circulating genotype over time.

Antibiotic utilisation data were extracted from National Antimicrobial Utilisation Surveillance Program data for our institution for the period 1 January 2015–30 September 2021. Bed occupancy data were extracted from the institutional information management system for the period 1 July 2014–30 June 2021.

Over the 5 year period April 2015–March 2020, the percentage of Enterobacteriaceae sterile site isolates expressing ESBLs was stable in our hospital at $2.2 \pm 0.3\%$ (Fig. 1A), with the predominant genotype being *bla*_{CTX-M} (72–88% of isolates from 2015–2019). Following the institution of international travel restrictions in March 2020, the overall proportion of sterile site Enterobacteriaceae isolates with an ESBL phenotype increased to 3.3% in the subsequent 12-month period (Fig. 1A).

Throughout the study period, the proportion of Enterobacteriaceae sterile site isolates producing CTX-M remained steady ($1.7 \pm 0.2\%$ for 2015–2019), and this continued through 2020 (1.7%) and the first quarter of 2021 (1.9%) (Fig. 1B). The proportion of IMP producing isolates remained low throughout (<0.2%). There was a notable rise in 2020 of isolates phenotypically testing as ESBL producing but testing negative for CTX-M, IMP-4 and pan-IMP genes on in house PCR testing (Fig. 1B), accounting for the rise in overall ESBL prevalence. Due to the retrospective nature of this study, more detailed genotypic information was not available.

Over the 6 year study period, antibiotic utilisation within the hospital rose by 46%, from 292 defined daily doses (DDD)/1000 bed days in 2015 to 426 DDD/1000 bed days in 2021, with the most significant rise (by 27%) occurring between 2019 and 2021 (Fig. 2A). The same trend was

Fig. 1 (A) Proportion of sterile site Enterobacteriaceae isolates producing extended spectrum beta-lactamases (ESBLs) by quarter [green line, mean case numbers per quarter for period 1 April 2015–31 March 2020; blue line, case numbers per quarter over 12-month period after international travel ban (1 April 2020–31 March 2021)]. (B) Yearly breakdown of proportion of Enterobacteriaceae sterile site isolates expressing ESBLs by genotype.

Fig. 2 (A) Antibiotic utilisation data at our institution for the period January 2015–September 2021, presented by antibiotic class. (B) Bed occupancy data from our institution by year. DDD, defined daily dose.

observed in usage of third generation cephalosporins in particular. While the current study addresses community acquired infections and hence is not directly influenced by inpatient antibiotic pressures, these hospital antibiotic utilisation rates suggest an overall increase in antibiotic usage over the study period which is likely directionally consistent with outpatient prescribing patterns.

In line with COVID-19 restrictions in 2020–2021, health seeking behaviours have changed in most of the world, with marked variability between settings. In our institution, admission rates rose by 3.4% in the 2020/2021 financial year compared to the average of the 5 years prior (average daily bed occupancy of 547 patients compared to 529 ± 11 patients in the previous period) (Fig. 2B). This small rise in hospital admissions is unlikely to have substantively affected ESBL circulation patterns.

This is a single centre study with several limitations. The COVID-19 pandemic brought unprecedented societal disruptions and widespread changes to the health care environment and to health seeking behaviours, which represent

several confounders to the current analysis. Utilising denominators in the current ESBL analysis, in the form of total Enterobacteriaceae isolates from sterile sites, as well as inclusion of local bed occupancy and antibiotic usage data, enables the ESBL incidence analysis to be placed in a more appropriate context.

Overall, these data suggest that local circulation of ESBLs, rather than international introduction, plays a major role in incidence of invasive multi-drug resistant (MDR) Gram negative infections in a low incidence, high income urban setting, at least in the medium term (up to 12 months following return from international travel). The observed rise in incidence of ESBL invasive disease is notable in the context of marked international travel restrictions but also in the presence of significant local non-pharmacological factors, including new physical distancing measures, a focus on hand hygiene and a short period of stay-at-home orders which might have been expected to limit transmission. Furthermore, the increased prevalence of ESBL-producing Enterobacteriaceae testing negative on established CTX-M/pan-IMP

molecular assays in 2020 is interesting. Larger prospective genotypic studies will enable more firm conclusions on emergence of new mutations and/or genotypes in the local population.

Recent studies from Italy and Taiwan revealed disparate findings with regard to AMR prevalence during the COVID-19 pandemic^{11,12} with high rates of AMR attributed in one study to increased antibiotic utilisation in COVID-19 patients.¹² In contrast to the experience in Europe and Asia, rates of SARS-CoV-2 infection were low in Australia through 2020 and early 2021, so factors relating directly to COVID-19 disease were not at play in this setting. Nonetheless, net antibiotic utilisation data from this study demonstrate a broad trend toward increased use over the period 2015–2021, and we hypothesise this is a major contributor to overall rising rates of invasive ESBL disease.

The observed rise in ESBL invasive disease suggests that local factors are key contributors to MDR Enterobacteriaceae infection in Australia. Alongside net increases in antibiotic use, other contributors may include inappropriate prescribing practices in hospital, community and agricultural settings, or prolonged carriage of MDR organisms in the gastrointestinal tract with invasive disease manifestation months to years later. This study lends weight to the important argument that considered antimicrobial stewardship is as critical in high income settings as it is in low income countries with higher baseline AMR prevalence rates.

AMR remains a critical public health issue faced locally and internationally and is a major focus of public health and research efforts. Interestingly, this study reveals that international travel is likely not the major contributor to the current ESBL epidemic, at least when viewed with a medium term lens. Rather, AMR must be owned as a local health priority driven at least in part by factors such as suboptimal antimicrobial stewardship at a community level. Future large scale studies are welcomed exploring longer term implications of travel restrictions on locally acquired versus internationally acquired ESBL infection rates, as well as ongoing surveillance for emerging mutations or genotypes.

Acknowledgements: The authors are grateful to the staff of the Nepean Hospital microbiology laboratory for their technical assistance and Mr Robert Robinson for helpful discussions.

Conflicts of interest and sources of funding: The authors state that there are no conflicts of interest to disclose.

Annaleise R. Howard-Jones^{1,2}, Samiha Rifaath Anver³, Kirsten Tsang⁴, James Branley^{1,2,3}

¹NSW Health Pathology-Nepean, Nepean Hospital, Kingswood, NSW, Australia; ²Faculty of Medicine and Health, University of Sydney, Camperdown, NSW, Australia; ³Department of Infectious Diseases and Microbiology, Nepean Hospital, Kingswood, NSW, Australia; ⁴Department of Pharmacy, Nepean Hospital, Kingswood, NSW, Australia

Contact Dr James Branley.

E-mail: james.branley@health.nsw.gov.au

1. Bezabih YM, Sabiti W, Alamneh E, *et al.* The global prevalence and trend of human intestinal carriage of ESBL-producing *Escherichia coli* in the community. *J Antimicrob Chemother* 2021; 76: 22–9.
2. O'Neill J. *Tackling Drug-Resistant Infections Globally: Final Report and Recommendations. The Review on Antimicrobial Resistance. May 2016.* London: Review on Antimicrobial Resistance, 2016. https://amr-review.org/sites/default/files/160525_Final%20paper_with%20cover.pdf.
3. Broaders E, Gahan CG, Marchesi JR. Mobile genetic elements of the human gastrointestinal tract: potential for spread of antibiotic resistance genes. *Gut Microbe* 2013; 4: 271–80.
4. Australian Commission on Safety and Quality in Health Care (ACSQHC). *AURA 2017: Second Australian Report on Antimicrobial Use and Resistance in Human Health.* Sydney: ACSQHC, 2017.
5. Solé M, Pitart C, Oliveira I, *et al.* Extended spectrum β -lactamase-producing *Escherichia coli* faecal carriage in Spanish travellers returning from tropical and subtropical countries. *Clin Microbiol Infect* 2014; 20: O636–9.
6. Pires J, Kraemer JG, Kuenzli E, *et al.* Gut microbiota dynamics in travelers returning from India colonized with extended-spectrum cephalosporin-resistant Enterobacteriaceae: a longitudinal study. *Travel Med Infect Dis* 2019; 27: 72–80.
7. Australian Bureau of Statistics (ABS). *Overseas Arrivals and Departures, Australia (April 2021).* Canberra: ABS, 2021.
8. Lee Y, Kim YA, Kim D, *et al.* Risk factors of community-onset extended-spectrum β -lactamase-producing *Klebsiella pneumoniae* bacteraemia in South Korea using national health insurance claims data. *Int J Antimicrob Agents* 2019; 54: 723–7.
9. Rawat D, Nair D. Extended-spectrum β -lactamases in Gram negative bacteria. *J Glob Infect Dis* 2010; 2: 263–74.
10. Birkett CI, Ludlam HA, Woodford N, *et al.* Real-time TaqMan PCR for rapid detection and typing of genes encoding CTX-M extended-spectrum beta-lactamases. *J Med Microbiol* 2007; 56: 52–5.
11. Bentivegna E, Luciani M, Arcari L, Santino I, Simmaco M, Martelletti P. Reduction of multidrug-resistant (MDR) bacterial infections during the COVID-19 pandemic: a retrospective study. *Int J Environ Res Publ Health* 2021; 18: 1003.
12. Lai CC, Chen SY, Ko WC, Hsueh PR. Increased antimicrobial resistance during the COVID-19 pandemic. *Int J Antimicrob Agents* 2021; 57: 106324.

DOI: <https://doi.org/10.1016/j.pathol.2021.11.005>

Identification of *Mycobacterium abscessus* to subspecies level with Bruker MALDI Biotyper

To the Editor,

Mycobacterium abscessus is a rapid growing mycobacteria (RGM) which is further separated into three subspecies: *Mycobacterium abscessus* subsp. *abscessus*, *Mycobacterium abscessus* subsp. *massiliense*, and *Mycobacterium abscessus* subsp. *bolletii*. They can cause various pulmonary and non-pulmonary infections, particularly in vulnerable patients with other underlying medical conditions. Macrolides are key antibiotics used in treating these patients, and are given in combination with other companion antimicrobials.^{1,2} Resistance to macrolides correlates with poorer outcomes. The most common mechanism of resistance is due to the presence of functional *erm* (41) in *M. abscessus* subsp. *abscessus* and *M. abscessus* subsp. *bolletii* which confers inducible resistance to clarithromycin.^{3,4} *Mycobacterium abscessus* subsp. *massiliense* is conversely associated with better outcomes due to truncation of *erm* (41) rendering it non-functional. Chromosomal mutations in the 23S rRNA (*rrl*) gene may result in constitutive macrolide resistance, however this is relatively uncommon.³